

Loughborough University Institutional Repository

Regional-scale drivers of groundwater faunal distributions

This item was submitted to Loughborough University's Institutional Repository by the/an author.

Citation: JOHNS, T. ... et al, 2015. Regional-scale drivers of groundwater faunal distributions. *Freshwater Science*, 34 (1), pp. 316 - 328.

Additional Information:

- This article was accepted for publication in the journal, *Freshwater Science* [University of Chicago Press / © Society for Freshwater Science]. The definitive version is available at: <http://dx.doi.org/10.1086/678460>

Metadata Record: <https://dspace.lboro.ac.uk/2134/18380>

Version: Accepted for publication

Publisher: University of Chicago Press / © Society for Freshwater Science

Rights: This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Please cite the published version.

1 Running head: Groundwater faunal distributions

2

3 REGIONAL SCALE DRIVERS OF GROUNDWATER FAUNAL DISTRIBUTIONS

4

5 Tim Johns^{1,6}, J. Iwan Jones², Lee Knight³, Louise Maurice⁴, Paul Wood⁵, Anne
6 Robertson^{6*}

7 1. Environment Agency, Red Kite House, Howbery Park, Wallingford, Oxfordshire,
8 OX10 8BD, UK.

9 2. School of Biological and Chemical Sciences, Queen Mary University of London,
10 London E14NS

11 3. No. 1 The Linhay, North Kenwood Farm, Oxtou, Nr Kenton, Devon, EX6 8EX

12 4. British Geological Survey, Maclean Building, Crowmarsh Gifford, Wallingford,
13 Oxfordshire, OX10 8BB UK.

14 5. Centre for Hydrological and Ecosystems Science, Department of Geography,
15 Loughborough University, Loughborough, Leicestershire, LE11 3TU, UK

16 6* Department of Life Sciences, University of Roehampton, Holybourne Avenue,
17 London SW15 4JD, UK. T: +44 (0) 208 392 3456 E: a.robertson@roehampton.ac.uk

18 .*Author to whom correspondence should be addressed

19

20

21

22

23 *ABSTRACT*

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

Freshwater aquifers are a major source of drinking water; they also possess unique assemblages of organisms. However, little is known about the distributional drivers of obligate groundwater organisms at the regional scale. We examine the distribution and composition of stygobiont assemblages in a complex geological setting and explore the relationship between groundwater fauna, hydrogeology and water chemistry. In the study area we grouped similar geologies into five hydrogeological formations (hydro-units) within which habitats for groundwater fauna were broadly similar. We found that the occurrence of stygobionts differed significantly between hydro-units. Stygobionts were significantly less likely to be recorded in mudstone/siltstone and sandstone aquifers compared with carbonate rocks or with igneous/metamorphic rocks. Variance partitioning indicated that the hydro-units explained a greater proportion of the variance (7.52%) in the groundwater community than water chemistry (5.02%). However, much of the variation remained unexplained. The macrofaunal stygobiont species in our study area formed three groups: (1) *Niphargus glenniei* was recorded in a range of hydro-units but only in the west of the study area. (2) *Niphargus kochianus*, *Niphargus fontanus*, *Proasellus cavaticus* and *Crangonyx subterraneus* were predominately recorded in carbonate aquifers in the east of the study area. (3) *Niphargus aquilex* and *Microniphargus leruthi*, were found throughout the study area and in a range of hydro-units. We hypothesise that physical barriers exist that prevent some stygobiont taxa from colonizing apparently suitable geologies; the low permeability deposits dividing the western and eastern parts of the study area may partly explain the observed distributions.

49

50

51 *KEYWORDS*

52 Hydrogeology, Geology, British Isles, Devon, Dorset, Stygobionts, Stygophiles,

53 *Niphargus*, Water chemistry, groundwater.

54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78

INTRODUCTION

Freshwater aquifers represent approximately 65% of drinking water sources in the EU (Scheidleder and Visser, 2012), and thus their hydrogeology and chemistry are generally well known. In addition to their value as a resource, groundwaters possess unique assemblages of organisms and globally harbour a large reservoir of biodiversity with a high proportion of endemic and rare species compared with surface water habitats (Sket 2004, Gibert et al. 2009). In response to the energy poor nature of subterranean environments, groundwater invertebrates typically have low population densities and extended life histories with delayed maturity, greater longevity, smaller egg-clutch size, larger eggs and lower percentages of mature ovigerous females than related surface-dwelling (epigean) species (Gibert et al. 1994). The low population numbers and slow reproduction of groundwater organisms may result in an increased vulnerability to anthropogenic impacts compared to epigean taxa. This vulnerability may be compounded by their slow dispersal rates (Culver et al. 2009).

Although research centred on surface-groundwater interactions is burgeoning and has moved from descriptive studies to a more experimental approach (introduction to this special issue, Larned 2012), research into deeper groundwaters is less developed, perhaps because of the difficulties in accessing this environment. Despite numerous studies describing the occurrence and distribution of groundwater fauna (e.g. Botosaneanu 1986, Proudlove et al. 2003, Robertson et al 2009, Gibert et al. 2009), relatively little is known about the environmental and/or historical drivers of their distribution patterns. The relative importance of a given driver is likely to vary across spatial and temporal scales. At the continental scale, stygobiont distribution

79 patterns have been shaped by events in deep geological time, for example the
80 movement of continental plates (e.g. Schminke 1974), orogenic and eustatic events
81 (Stock 1980, Boutin and Coineau 1990, Notenboom 1991), and the alternation of
82 marine transgressions and regressions (e.g. Jaume and Humphreys 2001). Long
83 term climate change has also shaped the distribution of groundwater taxa. In the
84 northern hemisphere the Pleistocene glaciations appear to have had a catastrophic
85 effect on groundwater communities, leaving Northern Europe with a depauperate
86 groundwater fauna compared to the rest of Europe. For example, only eight species
87 of obligate groundwater macro Crustacea have been recorded from mainland Britain,
88 including four species of *Niphargus*, the most speciose groundwater genus
89 (Proudlove et al. 2003, Robertson et al. 2009, Knight and Gledhill 2010). This
90 contrasts with 134 species of *Niphargus* recorded in the Balkans biogeographical
91 area (McInerney et al. 2014). In Northern Europe, contemporary distributions reflect
92 patterns of past glacial activity (e.g. Strayer 1994, Robertson et al. 2009). However,
93 there are some species present in formerly deglaciated areas, and those species
94 that remain in Northern Europe must have survived the impacts of glacial and
95 periglacial conditions (Holsinger 1988, Proudlove et al. 2003, Kristjansson and
96 Svavarsson 2007, Lefebure et al. 2007, McInerney et al. 2014). Indeed, recent
97 research has shown that the small number of *Niphargus* species that are present in
98 the UK are ancient endemics that have survived millions of years of extreme climate
99 changes (McInerney et al. 2014). At smaller scales, stygobiont distribution patterns
100 are influenced by the availability of suitable habitat and aquifer connectivity as well
101 as the ecological tolerances, competitive interactions and dispersal abilities of
102 species (Gibert et al. 1994, Castellarini et al. 2007, Hahn and Fuchs 2009).

104 Regional investigations that have gathered large datasets on stygobiont presence
105 and environmental conditions are relatively rare (Mauclaire and Gibert 2001, Galassi
106 et al. 2009, Martin et al. 2009, Dole-Olivier et al. 2009b, Hahn and Fuchs 2009). In
107 these studies, geology appears to be an important determinant of stygobiont
108 distributions because it controls the availability of suitable habitats (Datry et al. 2005,
109 Dole-Olivier et al. 2009b, Botosaneanu 1986, Hahn and Fuchs 2009). Most
110 stygobionts are dependent on well-developed void spaces with wide pore throats or
111 fracture apertures. These features commonly occur in karstic, porous and fractured
112 aquifers and this is where obligate groundwater communities are most diverse and
113 abundant (e.g. Gibert and Deharveng 2002, Dole-Olivier et al. 2009b). In addition to
114 providing physical habitat, the degree to which void spaces are developed and
115 connected also influences the exchange of oxygen, nutrients and organic matter with
116 the surface, and thus the availability of food for the groundwater community.

117 Observed distribution patterns may also relate to the opportunities for dispersal. In
118 groundwaters the habitat is very fragmented, limiting the dispersal of groundwater
119 fauna (Lefebure et al. 2007). Stygobiont assemblages are therefore characterised
120 by endemic species with limited geographical ranges. Even species which are
121 apparently widely distributed have emerged as cryptic lineages (Lefebure et al. 2007,
122 Trontelj et al. 2009, McInerney et al. 2014). Habitat fragmentation in groundwaters
123 has been most frequently discussed in relation to karst habitats (Trontelj et al. 2009,
124 Modovan et al. 2012), however it may also occur as a result of the juxtaposition of
125 different geologies. The presence of mudstones with low permeability and restricted
126 void spaces may form a physical barrier to stygobiont dispersal (Hahn and Fuchs
127 2009). Some stygobiont taxa, however, may be able to exploit superficial deposits or
128 the hyporheic zones of rivers as a means of dispersal (e.g. Ward and Palmer 1994,

129 Castellarini et al. 2007). The water chemistry of an aquifer, e.g. concentrations of
130 dissolved organic carbon and nitrates, may also correlate with the distribution of
131 groundwater taxa (Martin et al. 2009).

132

133 In this study we address the limited understanding of the drivers of groundwater
134 faunal distributions at the regional scale by carrying out an extensive sampling
135 programme within the complex geological setting of the south-west of the UK. Our
136 aims were to determine the detailed spatial distributions of the UK stygobiont species
137 in the study area and investigate whether the distribution and composition of
138 stygobiont assemblages varies with hydrogeology and water chemistry. We build on
139 the design of the PASCALIS project on European groundwater biodiversity which
140 adopted a hierarchical sampling strategy and included two types of aquifer (porous
141 and karstic) (Dole-Olivier et al. 2009a). Our approach was to group the different
142 hydrogeological formations present in the study area into five 'hydro-units' where the
143 different geologies within a given hydro-unit provide broadly similar habitats for
144 groundwater fauna. We sampled groundwater assemblages within the different
145 hydro-units to determine whether species were restricted to particular hydro-units,
146 and sampled water chemistry to determine whether faunal distributions reflect local
147 variation in water chemistry.

148

149

METHODS

Study area and hydrogeological units

151 The study area in south west England is an area of contrasting topography and
152 geology dominated by lowland (max. elevation 279 m) sedimentary rocks in the east
153 and higher elevation (max. 621 m) metamorphic and igneous rocks in the west (Fig.

154 1). The region's annual average rainfall is approximately 900-1000 mm
155 (<http://www.metoffice.gov.uk/climate/uk/sw/>).

156

157 Within the study area 271 different geological formations have been mapped at the
158 1:50,000 scale (Reproduced from the British Geological Survey Map data at the
159 original scale of 1:50 000 Licence 2011/057 British Geological Survey. ©NERC. All
160 rights reserved). To assess the general hydrogeological controls on species
161 distributions, these geologies were grouped into categories termed "hydro-units".
162 The hydro-units are based on broad differences in geology and groundwater flow
163 characteristics which result in different groundwater habitats for fauna. There are
164 two main types of groundwater flow: intergranular and flow through fractures (Price
165 1996). Unconsolidated granular deposits in which all groundwater movement is
166 intergranular were grouped together as hydro-unit 1 because these deposits form a
167 habitat in which fauna can only live within the pore spaces between the sediment
168 grains. Fractured rocks may be igneous, metamorphic or sedimentary in origin.
169 Igneous and metamorphic rocks (which are of volcanic origin) were grouped together
170 (hydro-unit 2) because there were insufficient data to determine differences in habitat
171 type between them. They were grouped separately from other strata because
172 groundwater flow is only through fractures, and due to of their volcanic origin, they
173 have a bedrock geochemistry that is different to other consolidated rocks.
174 Sedimentary rocks may be non-carbonate or carbonate. Non-carbonate
175 sedimentary rocks often have varying degrees of consolidation and, therefore, form
176 aquifers in which flow may be both inter-granular in the less consolidated areas and
177 through fractures in consolidated rock. They are classified according to grain size as
178 mudstones, siltstones and sandstones (Price 1996). Sandstones form aquifers with

179 much higher permeability than siltstones and mudstones which often form aquitards
180 or aquicludes (Price 1996, Allen et al. 1997). The difference in permeability is likely
181 to result in both a different groundwater chemistry and a different physical habitat for
182 fauna. They were therefore separated into a mudstone/siltstone hydro-unit (3) and a
183 sandstone hydro-unit (4). Hydro-unit 5 includes all carbonate geologies. In these
184 rocks, fractures are enlarged by dissolution to form fissures and caves providing
185 substantially larger and more inter-connected void spaces than any other rock type,
186 which can serve as habitats for fauna. In the study area carbonate rocks range from
187 the weakly karstified Cretaceous Chalk (Maurice et al, 2006) to the highly karstified
188 limestones of Devonian and Carboniferous. Where the Chalk overlies the Upper
189 Greensand Formation these strata are commonly in hydraulic continuity (Allen et al.
190 1997). This means that it is likely that fauna may be living in both the Chalk and the
191 Upper Greensand, and that it is not possible to distinguish the two habitats. The
192 Upper Greensand Formation is a non-karstic aquifer comprising sandstones and
193 sands, but where it is in hydraulic continuity with the Chalk it was included in the
194 carbonate hydro-unit (5). Elsewhere, outcrops of the Upper Greensand Formation
195 are isolated from the Chalk and, were therefore grouped with the other sandstones in
196 the sandstone hydro-unit (4). The properties of the five hydro-units are summarized
197 in Table 1.

198

199 The distribution and extent of the different hydro-units varied across the study area
200 (Fig. 1). In particular there was a pronounced difference between the hydro-units in
201 the east and west of the study area. Mudstones/siltstones (hydro-unit 3), covered the
202 largest surface area in the study region and were most common in the west.

203 Sandstones (hydro-unit 4), were the next most common, and occurred across the

204 region, with the largest outcrops in the west and north-west. Carbonate aquifers
205 (hydro-unit 5), were predominately located in the east, where there were extensive
206 Cretaceous chalk downlands. In the west carbonate rocks were limited to much
207 smaller isolated outcrops of older limestones, separated by mudstones and
208 sandstones. Granular material (hydro-unit 1) was restricted to the far south eastern
209 edge of the study area, whereas igneous and metamorphic rocks (hydro-unit 2),
210 were only present in the west and south west.

211

212 *Sampling*

213 In total, 221 sites were sampled. The number of samples taken within each hydro-
214 unit was approximately proportionate to the surface area it occupied (Table 1). Within
215 each unit, sites were selected to maximize spatial coverage. Boreholes free from fine
216 mesh well screens and in-situ equipment were preferentially selected. Borehole and
217 well sampling sites ranged from 0.36 to 130 m deep and 0.05 to 2 m in diameter.
218 Where there were insufficient boreholes or wells, springs were sampled. Each site
219 was sampled on a single occasion. Most samples (197) were collected between
220 September 2009 and May 2010, and water chemistry measurements were made at
221 the time of sampling. A further 24 samples were collected in November 2011; no
222 water chemistry data were obtained for these samples.

223

224 Fauna were collected from boreholes and wells using a weighted plankton net (63µm
225 mesh, diameter varied from 35 – 300mm dependent on borehole aperture). The net
226 was lowered to the base of the borehole/well and moved up and down vigorously to
227 disturb the sediment before retrieval. For springs, a net was swept through the
228 sediment of the spring chamber or at the spring head source. The samples were

229 preserved in >90% ethanol and all equipment thoroughly cleaned between sites. In
230 the laboratory animals were separated from debris, stygobionts were identified to
231 species level using Gledhill et al. (1993) and Knight and Gledhill (2010); Ostracoda
232 (identified to class), Hydracarina (identified to clade) and Copepoda and Oligochaeta
233 (identified to sub-class) were probably predominantly stygophiles and stygoxenes.
234 Freshwater epigeal taxa and terrestrial taxa were noted.
235 Electrical conductivity, pH, dissolved oxygen and temperature were recorded using a
236 multi-parameter probe (YSI 600QS) lowered to the base of each borehole/well. The
237 probe was calibrated prior to each sampling trip. Water samples for laboratory
238 analysis were taken using a bailer. At springs, field chemistry and water samples
239 were taken directly from the spring head or catchpit. Water samples were filtered and
240 refrigerated before dispatch to analytical laboratories within 48 hours. Analysis for
241 major anions (Cl^- , SO_4^{2-} , NO_3^-) and cations (Mg^{2+} , K^+ , Na^+ , Ca^{2+}) was performed
242 using high pressure liquid chromatography using a Dionex ED40 electrochemical
243 detector (min. reporting value; 1.0mg l^{-1}). Ammoniacal nitrogen was analyzed by
244 colourimetric analysis using a Konelab discrete analyser (min. reporting value; 0.03
245 mg l^{-1}). Dissolved organic carbon (DOC) was analyzed by a Skalar San Plus 1050 air
246 segmented flow analyser (min. reporting value 0.2 mg l^{-1}).

247

248 *Data analysis*

249 Chemical data were tested for normality (Q-Q plots) and homogeneity of variance
250 (Fligner-Killeen test), and a non-parametric ANOVA was used to investigate
251 differences in water chemistry between the five different hydro-units (selected as
252 factors). A generalized linear model with a binomial error structure was then used to
253 examine the relationships between presence/absence of stygobiont taxa, hydro-units

254 and water chemistry. Statistical analyses were performed using R version 3.0.2
255 (2013), using standard packages plus the package for analysis of over dispersed
256 data (Lesnoff et al, 2012). The initial model selected hydro-unit as a factor. The
257 model was then run with all stygobiont presence/absence data, using the carbonate
258 hydro-unit (5) as the point of comparison because it: i) comprised the most sites, ii)
259 was the strata in which stygobionts were most common, and iii) also yielded a broad
260 range of water quality data. Models were then run for each set of water quality data
261 (i.e. each chemical determinand with associated presence/absence and hydro-unit
262 data), the odds ratios were calculated and predicted probabilities of stygobiont
263 presence were determined from the mean value for each determinand. Different
264 permutations of the model were run to explore whether the inclusion of sample type
265 (borehole, well or spring) and/or water quality data improved the model, and to
266 identify any interactions between factors. Outputs were checked to verify the data
267 were not over dispersed and models were then compared, using a one-way ANOVA,
268 to identify any significant differences between models. Where there was no
269 difference, the simplest model was selected and the odds ratios calculated to
270 describe the relative likelihood of recording stygobionts in each hydro-unit.

271

272 The association between the faunal community and environmental variables was
273 investigated using Canonical Correspondence Analysis (CCA) within Canoco for
274 Windows Ver.4.51 (Ter Braak and Smilauer 1998). Only those sites with concurrent
275 chemistry analysis (190 samples) were used in this analysis and sites where there
276 was uncertainty over the geological unit from which the sample was obtained (7
277 samples) were excluded.

278 A series of ordinations were undertaken including groundwater faunal abundance,
279 hydro-units and water chemistry data. Both stygobiont and stygophile taxa were
280 included in the analysis to reflect the diversity and distribution of the community
281 recorded from each sample location where fauna were recorded, although the
282 results and discussion are focused on the stygobiont taxa. To determine how
283 individual variables might explain patterns in the faunal data (stygobionts and
284 stygophiles), we applied the forward selection procedure ($P \leq 0.05$ after Bonferonni
285 correction). Only those variables having a significant association with the faunal
286 community distribution were included in the final analysis and presented in the output
287 diagrams. The stygobiont *Antrobathynella stammeri* (Jakobi, 1954), which was
288 recorded only once, was fitted passively to the ordination. To determine the
289 proportion of variance in the faunal data explained by the hydro-units and water
290 chemistry data, a variance partitioning approach was used (Borcard 1992). This
291 constrained the species ordination for each of these co-variable groups in turn, while
292 partialling out the other variables from the ordination.

293

294

295

RESULTS

296 As expected water chemistry varied significantly between the five hydro-units,
297 reflecting differences in bedrock geochemistry and permeability. With the exception
298 of potassium, significant differences were recorded between each chemical
299 determinand and the five hydro-units (Table 2).

300

301 Stygobionts and/or stygophiles were recorded from 70% of the study sites (Table 3)
302 and all hydro-units (Fig.1). The remaining 30% of sites contained no groundwater

303 taxa although epigean taxa were recorded at some sites. Stygophiles were more
304 abundant and recorded more frequently than stygobionts (59% compared to 34% of
305 sites). All of the stygobiont macro-Crustacea known to occur in the UK were
306 recorded, although their presence and community composition varied considerably
307 across the study region. Boreholes in the carbonate hydro-unit (unit 5) supported the
308 most diverse stygobiont communities, with up to four taxa recorded in a single
309 borehole, compared to a maximum of two taxa from other hydro-units (Fig. 2).

310

311 *Niphargus aquilex* (Schiödte, 1855), was recorded throughout the study region and
312 from all hydro-units (Fig. 2). In contrast, *Niphargus glenniei* (Spooner, 1952) was
313 only recorded in the west of the study area despite being recorded in most hydro-
314 units. *Niphargus kochianus* (Bate, 1859), *Niphargus fontanus* (Bate, 1859) and
315 *Proasellus cavaticus* (Leydig, 1871) were primarily recorded in the carbonate hydro-
316 unit (5) in the east of the region (Fig. 2), but were absent from this hydro-unit and all
317 other hydro-units in the west of the study area. *Crangonyx subterraneus* (Bate,
318 1859) and *Microniphargus leruthi* (Schellenberg 1934) were rare, and were primarily
319 recorded in the east (Fig. 2). *Antrobathynella stammeri* (Jakobi, 1954) was recorded
320 once.

321

322 The occurrence of stygobionts differed significantly between hydro-units [$\chi^2=33.8$,
323 $df=5$, $p<0.001$, Table 4]. From the odds ratio (OR), stygobionts were 88% less likely
324 to be recorded in mudstone/siltstones (hydro-unit 3) [OR=0.12, 97.5%CI 0.034-
325 0.030, $p<0.001$] than in carbonates (hydro-unit 5) and about 80% less likely to be
326 recorded in sandstones (hydro-unit 4) [OR=0.20, 97.5%CI 0.08-0.50, $p<0.001$], than
327 carbonates (hydro-unit 5). Stygobionts were also less likely to be recorded in

328 mudstones/siltstones (hydro-unit 3) and sandstones (hydro-unit 4) than in
329 igneous/metamorphic sites (hydro-unit 2) and granular sites (hydro-unit 1). All
330 differences were significant except for sandstone sites relative to granular sites.
331 The generalized linear model generally improved (based on the AIC scores and level
332 of significance) with the addition of either sample type or a water chemistry
333 determinand as a value. Mudstones/siltstones and sandstones (hydro-units 3 and 4)
334 were significantly different from carbonate rocks (hydro-unit 5) throughout the
335 iterations. No significant interactions between factors (e.g. hydro-unit combined with
336 sample type) were identified. By including water chemistry determinands with
337 presence/absence in the model, improved models were obtained with calcium,
338 chloride and pH , indicating a potential association of these variables with stygobiont
339 presence. The predicted probability of stygobiont presence in any hydro-unit was
340 higher with increasing calcium concentrations. In contrast, the predicted probability
341 of stygobiont presence in any hydro-unit declined as chloride concentrations
342 increased. There was no clear relationship between pH and the presence of
343 stygobionts.

344

345 The inclusion of sample type significantly improved the models that included
346 electrical conductivity, magnesium, potassium, sodium, sulphate and dissolved
347 organic carbon. For the model with potassium, the igneous/metamorphic aquifers
348 (hydro-unit 2) were identified as significantly different when sample type was
349 included. The odds ratio of recording stygobionts in this unit was also lower
350 compared with the simplest model where no additional factors were included. Overall
351 there was a greater likelihood of recording stygobionts in springs (sample type 3)
352 than boreholes (sample type 1: Table 4). However, presence or absence of

353 stygobionts across the hydro-units was significant ($p < 0.001$) irrespective of the
354 inclusion of sample type in the model.
355
356 Canonical correspondence analysis indicated a separation of the sample centroids
357 for the five hydro-units based on the faunal community data (Fig. 3), the first two
358 canonical axes explained 20.0% and 13.6% of the total variance in faunal
359 abundance respectively (Fig. 3). These two axes combined explained 78.4% of the
360 variance in the species-environment relationship (46.7% and 31.7% respectively).
361 Water temperature ($^{\circ}\text{C}$), magnesium (mg/l) and potassium (mg/l) were significantly
362 correlated with the distribution of the groundwater fauna (all $p < 0.01$). The stygobionts
363 *Niphargus kochianus*, *Microniphargus leruthi*, *Crangonyx subterraneus* and
364 *Proasellus cavaticus* were associated with the centroids for the sandstone and
365 carbonate hydro-units (hydro-unit 4 and 5). In contrast, some taxa were not strongly
366 associated with the centroids of any single hydro-unit but were associated with
367 higher concentrations of potassium (*N. fontanus* and *N. aquilex*) and higher water
368 temperature (*N. glenniei*). Variance partitioning indicated that 7.5% of the variance in
369 faunal abundance could be explained by the hydro-units and 5.0% by water
370 chemistry (Table 5), whilst 3.6% of the variance was shared by hydro-units and water
371 chemistry together.

372

373

DISCUSSION

374 Stygobionts and stygophiles were found throughout the region and were present in
375 all hydro-units. However, stygobionts were absent from 70% of sites. Eberhard et al.
376 (2009) and Hancock and Boulton (2009) found that repeated sampling over time at
377 groundwater sites increased the number of species recorded, thus we cannot

378 exclude the possibility that the low frequency of stygobiont occurrence in our study
379 may be an artifact of low sampling effort, as we sampled each site on only one
380 occasion.

381 We found that stygobiont occurrence and community composition differed between
382 hydro-units. Stygobionts were significantly more likely to be recorded in the
383 carbonate and igneous/metamorphic hydro-units, than in the mudstone/siltstone or
384 sandstone units. Several stygobiont species (*Niphargus kochianus*, *Microniphargus*
385 *leruthi*, *Crangonyx subterraneus* and *Proasellus cavaticus*) were predominantly
386 associated with the carbonate hydro-unit. Other stygobionts (*Niphargus aquilex* and
387 *Niphargus glenniei*) were found across a range of hydro-units.

388 The carbonate hydro-unit also supported the most diverse stygobiont communities,
389 with up to four stygobiont species co-occurring in a single borehole. Carbonate
390 aquifers are characterised by solutional fissures and conduits providing connected
391 water-filled void spaces which are sufficiently large for groundwater organisms to
392 inhabit (Humphreys 2009). Furthermore, their highly permeable nature may result in
393 a greater supply of food from the surface, permitting the co-existence of more
394 species than in less permeable hydro-units, where food shortages may be severe
395 (Malard et al. 2009). Studies in continental Europe have demonstrated previously
396 that karst aquifers support some of the most diverse stygobiont communities (Dole-
397 Olivier et al., 2009b; Martin et al., 2009; Galassi et al., 2009; Hahn and Fuchs, 2009).

398 The sandstone hydro-unit was also anticipated to provide a suitable habitat for
399 diverse stygobiont communities because this has been the case in other regions
400 (Humphreys 2009). Therefore, the low numbers of individuals and low diversity
401 recorded in our study was surprising. One explanation is that the Permo-Triassic

402 sandstones, which comprise part of this hydro-unit contain lower permeability
403 mudstone beds in this area (Allen et al. 1997). This may result in a fragmented
404 habitat with areas containing poorly developed fracture networks providing a poor
405 habitat. The deposits are also affected by faulting which may result in further
406 barriers to stygobiont movement. However, the sandstone (hydro-unit 4) is
407 predominantly distributed in the west of the study area, which is outside the
408 observed range of *Niphargus kochianus*, *Niphargus fontanus* and *Proasellus*
409 *cavaticus* (Fig. 2), three of the most common stygobite species in the east of the
410 study area. It is not clear whether the absence of these species in the west is
411 because the sandstones do not form a suitable habitat for them, or because they
412 have not dispersed to these areas.

413 The mudstone/siltstone hydro-unit had the lowest occurrence of stygobionts
414 suggesting that it provides a less suitable habitat, perhaps as a result of the small
415 and often weakly connected voids, and the low levels of oxygen and nutrients. Our
416 findings are consistent with those of Hahn and Fuchs (2009), who also reported that
417 mudstones were characterized by depauperate communities.

418

419 Our results demonstrate that the five hydro-units had different water chemistries.
420 Therefore it was not possible to determine whether differences in groundwater fauna
421 occurred due to water chemistry differences between the units, or differences
422 between the physical voids present within the different units (Maurice and
423 Bloomfield, 2012). However, in this study, groundwater taxa occurred in all units and
424 the proportion of the variance in the abundance of the groundwater community
425 explained by water chemistry was smaller than that of the hydro-units. The
426 mudstone/siltstone and sandstone units remained significantly different from the

427 carbonate unit in the general linear model, irrespective of the addition of water
428 chemistry parameters. However, the igneous/ metamorphic unit only became
429 significantly separated from the other hydro units when potassium (together with
430 sample type) was included in the model. The higher potassium concentrations
431 reported from the igneous/ metamorphic hydro-unit relative to the carbonate hydro-
432 unit reflects the contrasting water chemistry of these different geologies. These
433 findings provide evidence that suggests that, across our study sites, water chemistry
434 had less influence over the distribution of groundwater taxa than physical
435 hydrogeological characteristics. Other studies have also found little relation between
436 environmental variables, water chemistry and the distribution of groundwater taxa at
437 a regional scale (e.g. Dumas and Lescher-Moutoue 2001, Paran et al. 2005, Dole-
438 Olivier et al. 2009b, Galassi et al. 2009). Nevertheless, in all these studies there are
439 considerable difficulties when trying to separate the influence of potentially
440 confounding factors that occur over a range of geographical scales that may
441 determine stygobiont distributions (Dole-Olivier et al., 2009b; Stoch and Galassi,
442 2010; Maurice and Bloomfield, 2012).

443

444 The probability of finding stygobionts appeared to increase with an increase in
445 calcium concentrations. This is probably because stygobionts were most often found
446 in the carbonate hydro-unit where calcium concentrations are high. Although
447 calcium is necessary for crustacean carapace development (Rukke, 2002), it does
448 not appear to be limiting in our study area as high stygobiont frequencies were also
449 recorded from the igneous/metamorphic hydro-unit, where the mean calcium
450 concentration was the lowest of all the hydro-units sampled.

451

452 In contrast to Dole-Olivier et al. (2009b), we found no significant relationship
453 between dissolved oxygen concentration and stygobiont occurrence. This is
454 probably because all but 14 of our sites had oxygen concentrations $>1 \text{ mg l}^{-1}$, which
455 is the threshold below which oxygen has been reported to be critical for groundwater
456 taxa (Malard and Hervant 1999, Hahn 2006). In addition, stygobionts are widely
457 acknowledged to be tolerant of low oxygen conditions in comparison to epigean
458 species (Malard and Hervant 1999).

459 The greater likelihood of finding stygobionts in spring sites (sample type 3) than in
460 borehole sites (sample type 1) may be because it is easier to capture fauna in
461 springs than in boreholes, and also because springs form a distinct ecotone where
462 there are higher nutrients and more diverse communities than in deeper
463 groundwaters (Smith et al. 2003) However, the use of springs as an explanatory
464 variable did not fundamentally modify or significantly influence the differences we
465 identified between the hydro-units.

466

467 Most of the variation in the occurrence of groundwater fauna across the study region
468 was not explained by either the hydro-units or water chemistry. One of our most
469 striking findings was the differing distribution patterns of stygobiont species within the
470 study area. Most stygobiont species had restricted, and different distributions. The
471 exception was *Niphargus aquilex* which was recorded from all hydro-units and
472 across the whole study area. This species is widely distributed in central and
473 southern Europe also (Botosaneanu 1986), with an apparent range that spans $>$
474 2300 km east – west (Trontelj et al. 2009). In the British Isles it has been extensively
475 recorded in riverine hyporheic zones and is particularly associated with shallow
476 alluvial aquifers (Proudlove et al. 2003) and, thus, may have dispersed across the

477 area using this route (the hyporheic corridor concept of Stanford and Ward 1993).
478 Nevertheless, recent research shows that the wide distribution of *Niphargus aquilex*
479 is an artifact generated by cryptic diversity; the complex contains seven cryptic taxa,
480 two of which are unique to the British Isles (McInerney et al. 2014). Other widely
481 distributed *Niphargus* “species” have also been found to be species complexes (e.g.
482 *N. virei*, Lefebure et al. 2006 and *N. rhenorhodanensis*, Lefebure et al 2007, see also
483 Trontelj et al. 2009).

484

485 Although there would appear to be suitable habitats distributed across the study
486 area, other stygobiont species exhibited distinct spatial segregation. The endemic
487 *Niphargus glenniei* was only recorded in the west and *Niphargus kochianus*,
488 *Niphargus fontanus*, and *Proasellus cavaticus* were only found in the east. One
489 possible explanation for these observed distributions is that they result from biotic
490 interactions between the species. *Niphargus glenniei* for example, did not co-occur
491 with either *Niphargus fontanus* or *Niphargus kochianus*. *Niphargus fontanus* is much
492 larger and may have a competitive or predatory advantage. However, stygobiont
493 species can co-exist, and up to four species were found in a single borehole in the
494 east of our study area. Although in our study *Niphargus glenniei* was mostly
495 recorded alone or occasionally together with the smaller species *Microniphargus*
496 *leruthi*, other studies have reported it with the larger species *Niphargus aquilex*
497 (<http://hcrs.freshwaterlife.org/>). An alternative hypothesis is that geological barriers
498 are present that constrain the dispersal of these organisms, as has been reported for
499 other groundwater taxa (Gooch and Hetrick 1979; Goricki and Trontelj 2006).
500 Trontelj et al. (2009) observed that many groundwater taxa appear to be able to
501 disperse and yet are confined by boundaries beyond which dispersal is impossible

502 (although the nature of these boundaries is currently unclear). The
503 mudstones/siltstones and other low permeability rocks across the centre of our study
504 area may form a barrier between suitable stygobiont habitats in the east and west,
505 and could explain the different species distributions. However, given the occurrence
506 of all stygobiont species in spring habitats, it is surprising that groundwater species
507 have not been able to utilize shallow superficial deposits (e.g. hyporheic zone
508 corridors) to disperse between the east and west of the study area.

509 The life histories and restricted distributions of groundwater taxa mean that they are
510 vulnerable to anthropogenic change (Gibert et al. 1994). *Niphargus glenniei*, which is
511 endemic to two counties in southern England, has been designated a UK
512 Biodiversity Action Plan species. However, other groundwater taxa have no such
513 recognition, and European groundwater monitoring programmes do not consider
514 groundwater ecology. Identifying the drivers of observed distribution patterns in
515 groundwater taxa is important because this information may then be used to inform
516 the development of management plans for the conservation of these unique
517 assemblages.

518 .

519 CONCLUSIONS

520 In our regional scale study, there were significant differences in the groundwater
521 assemblages present in geologies with different hydrogeological characteristics.
522 Stygobionts occurred most frequently and were most diverse in the carbonate hydro-
523 unit (5) which is characterized by karstic, solutionally enlarged voids providing an
524 extensive physical habitat, and a good supply of nutrients and oxygen. They were
525 also relatively abundant and diverse in the igneous and metamorphic hydro-unit (2)
526 and granular aquifers (hydro-unit 1) but were relatively rare in the sandstone hydro-

527 unit (4). They occurred least frequently in the mudstone/ siltstone hydro-unit (3)
528 which has low permeability resulting in poor physical and water chemistry conditions
529 for stygobionts. The hydro-units explained a greater proportion of the variation in
530 stygobiont data than water chemistry, suggesting that physical hydrogeology may be
531 more important than water chemistry in determining stygobiont distributions.
532 However, a large proportion of the variation in stygobiont abundance was not related
533 to either the hydro-units or water chemistry and we were intrigued to find that some
534 stygobiont species were spatially segregated within our study area. It appears that
535 barriers may exist that prevent some stygobiont taxa from colonizing apparently
536 suitable geologies, and the low permeability deposits dividing the western and
537 eastern parts of the study area may partly explain the observed distributions. The
538 stygobiont species fell into three contrasting groups: (1) *Niphargus glenniei* was only
539 found in the west of the study area, but was found in a range of hydro-units with
540 varying void types and water chemistry. (2) *Niphargus kochianus*, *Niphargus*
541 *fontanus*, *Crangonyx subterraneus* and *Proasellus cavaticus* were only found in the
542 east of the study area and predominantly in carbonate aquifers. (3) *Niphargus*
543 *aquilex* and *Microniphargus leruthi*, were found throughout the study area and in a
544 range of hydro-units suggesting that they have good dispersal abilities and may be
545 able to circumvent low permeability barriers, perhaps by utilizing superficial deposits
546 or hyporheic zone corridors. Identifying the precise nature of the barriers to dispersal
547 and the development of monitoring and conservation plans must be a priority for
548 these restricted and vulnerable species.

549

550

ACKNOWLEDGMENTS

551 This research was funded by the Esmee Fairbairn Foundation, grant number 08-
552 101,7 to ALR, PJW, LK and LM. We thank the many landowners who gave us
553 permission to cross their land, and the Environment Agency for their help and
554 assistance in accessing sites and provision of equipment. L. Maurice publishes with
555 the permission of the Executive Director, British Geological Survey (NERC).

556

557

LITERATURE CITED

- 558 Allen, D.J., L.J. Brewerton, L.M. Coleby, B.R. Gibbs, M.A. Lewis, A.M. MacDonald,
559 S.J. Wagstaff, and Williams, A.T., 1997. The physical properties of the major
560 aquifers in England and Wales. BGS Technical Report WD/97/34, Environment
561 Agency R&D Publication 8. 312 pp
- 562 Botosaneanu, L. 1986. *Stygofauna Mundi*. E.J. Brill, Leiden.
- 563 Borcard, D., P. Legendre, and P. Drapeau, 1992. Partialling out the spatial
564 component of ecological variation. *Ecology* 73:1045-1055.
- 565 Boutin, C., and N. Coineau. 1990. "Regression model, "modèle biphasé" d'évolution
566 et origine des microorganismes stygobies interstitiels continentaux. *Revue de*
567 *Micropaléontologie* 33:303–322
- 568 Castellarini, F., F. Malard, M-J. Dole-Olivier, and J. Gibert. 2007. Modelling the
569 distribution of stygobionts in the Jura Mountains (eastern France). Implications for
570 the protection of ground waters. *Diversity and Distributions* 13:213-224.
- 571 Culver D.C., T. Pipan, and K. Schneider. 2009. Vicariance, dispersal and scale in
572 the aquatic subterranean fauna of karst regions. *Freshwater Biology* 54:918-929.
- 573 Datry, T., F. Malard, and J. Gibert. 2005. Response of invertebrate assemblages to
574 increased groundwater recharge rates in a phreatic aquifer. *Journal of the North*
575 *American Benthological Society* 24:461-477.

576 Dole-Olivier, M.-J., F. Castellarini, N. Coineau, D.M.P. Galassi, P. Martin, N. Mori, A.
577 Valdecasas, and J.Gibert. 2009a. Towards an optimal sampling strategy to
578 assess groundwater biodiversity: comparison across six European regions.
579 *Freshwater Biology* 54:777-796.

580 Dole-Olivier, M.-J., F. Malard, D. Martin, T. Lefebure, and J. Gibert. 2009b.
581 Relationships between environmental variables and groundwater biodiversity at
582 the regional scale. *Freshwater Biology* 54:797-81

583 Dumas, P., and F. Lescher-Moutoue. 2001. Cyclopid distribution in an agriculturally
584 impacted alluvial aquifer. *Archiv für Hydrobiologie* 150:511-528.

585 Eberhard, S. M., S.A. Halse, M.R. Williams, M.D. Scanlon, J. Cocking, and H.J.
586 Barron. 2009. Exploring the relationship between sampling efficiency and short-
587 range endemism for groundwater fauna in the Pilbara region, Western Australia.
588 *Freshwater Biology* 54:885-901.

589 Galassi, D.M.P., F. Stoch, B. Fiasca, T. Di Lorenzo, and E. Gattone. 2009.
590 Groundwater biodiversity patterns in the Lessinian Massif of northern Italy.
591 *Freshwater Biology* 54:830-847.

592 Gibert, J., D. Danielopol, and J. Stanford (Eds). 1994. *Groundwater Ecology*.
593 Academic Press LTD, London UK. pp1-551.

594 Gibert, J., D.C. Culver, M.-J. Dole-Olivier, F. Malard, M.C. Christman, and L.
595 Deharveng. 2009. Assessing and conserving groundwater biodiversity: synthesis
596 and perspectives. *Freshwater Biology* 54:930-941.

597 Gibert, J., and L. Deharveng. 2002. Subterranean ecosystems: A truncated
598 functional biodiversity. *Bioscience* 52:473-481.

599 Gledhill, T., D.W. Sutcliffe, and W.D. Williams. 1993. *British Freshwater Crustacea*
600 *Malacostraca: A key with ecological notes*. Freshwater Biological Association

601 Gooch J.L., and S.W. Hetrick. 1979. The Relation of Genetic Structure to
602 Environmental Structure: *Gammarus minus* in a KarstArea. *Evolution* 33:192-206.

603 Goricki, S., and P. Trontelj. 2006. Structure and evolution of the mitochondrial control
604 region and flanking sequences in the European cave salamander *Proteus*
605 *anguinus*. *Gene* 378:31-41.

606 Hahn, H.J. 2006. The GW-Fauna-Index: A first approach to a quantitative ecological
607 assessment of groundwater habitats. *Limnologica* 36:119-137.

608 Hahn, H.J., and A. Fuchs, 2009. Distribution patterns of groundwater communities
609 across aquifer types in south-western Germany. *Freshwater Biology* 54:848-860.

610 Hancock, P.J., and A.J. Boulton. 2009. Sampling groundwater fauna: efficiency of
611 rapid assessment methods tested in bores in eastern Australia. *Freshwater*
612 *Biology* 54:902-917.

613 Holsinger J.R.1988. Troglobites: The evolution of cave dwellin organisms. *American*
614 *Scientist* 76:146-153.

615 Humphreys W.F. 2009. Hydrogeology and groundwater ecology: Does each inform
616 the other? *Hydrogeology Journal* 17:5-21

617 Jaume, D., and W.F. Humphreys. 2001. A new genus of epacteriscid calanoid
618 copepod from an anchialine sinkhole in northwestern Australia. *Journal of*
619 *Crustacean Biology* 21:157–169.

620 Knight, L.R.F.D. and T. Gledhill. 2010. The discovery of *Microniphargus leruthi*
621 Schellenberg, 1934 (Crustacea: Amphipoda: Niphargidae) in Britain and its
622 distribution in the British Isles. *Zootaxa* 2655:52-56.

623 Kristjansson B.K., and J. Svavarsson. 2007. Sub-glacial refugia in Iceland enabled
624 groundwater amphipods to survive glaciations. *American Naturalist* 170:292-296.

625 Larned S.T. 2012. Phreatic groundwater ecosystems: research frontiers for
626 freshwater ecology. *Freshwater Biology* 57:885-906.

627 Lefébure, T., C.J. Douady, M. Gouy, P. Trontelj, J. Briolay, and J. Gibert. 2006.
628 Phylogeography of a subterranean amphipod reveals cryptic diversity and
629 dynamic evolution in extreme environments. *Molecular Ecology* 15:1797–1806.

630 Lefébure, T., C.J. Douady, F. Malard, and J. Gibert. 2007. Testing dispersal and
631 cryptic diversity in a widely distributed groundwater amphipod (*Niphargus*
632 *rhenorhodanensis*) *Molecular Phylogenetics and Evolution* 42:676–686.

633 Lesnoff, M., Lancelot, R. (2012). aod: Analysis of Overdispersed Data. R package
634 version 1.3, URL <http://cran.r-project.org/package=aod>

635 Malard, F., and F. Hervant. 1999. Oxygen supply and the adaptations of animals in
636 groundwater. *Freshwater Biology* 41:1-30.

637 Malard, F., C. Boutin, A.I. Camacho, D. Ferreira, G. Michel, B. Sket, and F. Stoch.
638 2009. Diversity patterns of stygobiotic crustaceans across multiple scales in
639 Europe. *Freshwater Biology* 54: 756-776.

640 Martin, P., C. De Broyer, F. Fiers, G. Michel, R. Sablon, and K. Wouters. 2009.
641 Biodiversity of Belgian groundwater fauna in relation to environmental conditions.
642 *Freshwater Biology* 54:814-829.

643 Mauclaire, L. and J. Gibert. 2001. Environmental determinants of bacterial activity
644 and faunal assemblages in alluvial riverbank aquifers. *Archiv für Hydrobiologie*
645 152:469-487.

646 Maurice, L. and J. Bloomfield. 2012. Stygobitic invertebrates in groundwater — A
647 review from a hydrogeological perspective. *Freshwater Reviews* 5:51-71.

648 Maurice, L.D., T.C. Atkinson, J.A. Barker, J.P. Bloomfield, A.R. Farrant, and A.T.
649 Williams. 2006. Karstic behaviour of groundwater in the English Chalk. *Journal of*
650 *Hydrology*. 330: 63-70.

651 McInerney C.E., L. Maurice, A.L. Robertson, L.R.F.D. Knight, J. Arnsheidt, C.
652 Venditti, J.S.G. Dooley, T. Mathers, S. Matthijs , K. Eriksson, G. Proudlove and B.
653 Hänfling (2014) The ancient Britons: Groundwater fauna survived extreme
654 climate changes over tens of millions of years across NW Europe. *Molecular*
655 *Ecology* 23: 1153-1166.

656 Moldovan O.T., I.N. Meleg, and A. Persoiu. 2012. Habitat fragmentation and its
657 effect on groundwater populations. *Ecohydrology* 5:445-452.

658 Notenboom, J., 1991. Marine regressions and the evolution of groundwater dwelling
659 amphipods (crustacea). *Journal of Biogeography* 18:437-454.

660 Paran F., F. Malard J. Mathieu, M. Lafont, D.M. P. Galassi, and P. Marmonier. 2005.
661 Distribution of groundwater invertebrates along an environmental gradient in a
662 shallow water-table aquifer. Pages 99-105 in J. Gibert (editor) *World*
663 *Subterranean Biodiversity, Proceedings of an International Symposium* (Ed. J.
664 Gibert) University Claude Bernard, Lyon, France

665 Price M. 1996. *Introducing Groundwater*, 2nd edn. Chapman and Hall, London. pp
666 70-87.

667 Proudlove, G.S., P.J. Wood, P.T. Harding, D.J. Horne, T. Gledhill, and L.R.F.D.
668 Knight.2003. A review of the status and distribution of the subterranean aquatic
669 Crustacea of Britain and Ireland. *Cave and Karst Science* 30:51-74

670 R Core Team (2013). *R: A language and environment for statistical computing*.
671 R Foundation for Statistical Computing, Vienna, Austria. URL [http://www.R-proje](http://www.R-project.org/)
672 [ct.org/](http://www.R-project.org/).

673 Robertson, A. L., J.W.N. Smith, T. Johns, and G.S. Proudlove. 2009. The distribution
674 and diversity of stygobites in Great Britain: an analysis to inform groundwater
675 management. *Quarterly Journal of Engineering Geology and Hydrogeology*
676 42:359-368.

677 Rukke, N.A. 2002. Effects of low calcium concentration on two common freshwater
678 crustaceans, *Gammarus lacustris* and *Astacus astacus*. *Functional Ecology*
679 16:357–366.

680 Scheidleder, A., and P. Visser, 2012. Comparative Study of Pressures and
681 Measures in the Major River Basin Management Plans: Drinking water protected
682 areas: Task 3c – Summary note on drinking water safeguard zones. European
683 Commission. 3-4.

684 Schminke, H,K. 1974. Mesozoic Intercontinental Relationships as Evidenced by
685 Bathynellid Crustacea (Syncarida : Malacostraca). *Systematic Biology* 23:157-
686 164.

687 Sket, B. 2004. Biodiversity in hypogean waters page 709. *In: J. Gunn (editor)*
688 *Encyclopedia of Caves and Karst Science*. Fitzroy Dearborn, New York,

689 Smith, H., P.J. Wood, and J. Gunn 2003. The influence of habitat structure and flow
690 permanence on invertebrate communities in karst spring systems. *Hydrobiologia*
691 510: 53-66.

692 Stanford, J.A., and J.V. Ward. 1993. An ecosystem perspective of alluvial rivers;
693 connectivity and the hyporheic corridor. *Journal of the North American*
694 *Benthological Society* 12:48-60.

695 Stock, J., 1980. Regression model evolution as exemplified by the genus
696 *Pseudoniphargus* (Amphipoda). *Bijdragen tot de Dierkunde* 50:105–144.

697 Stoch, F., and D.M.P. Galassi, 2010. Stygobiotic crustacean species richness: a
698 question of numbers, a matter of scale. *Hydrobiologia* 653:217-234.

699 Strayer, D.L., 1994. Limits to biological distributions in groundwater. *in* J. Gibert, D.
700 Danielopol and J. Stanford. (editors) *Groundwater Ecology*. Academic Press
701 Limited, London. U.K. pp 287-310.

702 ter Braak, C.J.F and P. Smilauer. 1998. CANOCO reference manual and user's
703 guide to Canoco for windows: software for canonical community ordination
704 (version 4). Wageningen.

705 Trontelj, P., C.J. Douady, C. Fiser, J. Gibert, S. Goricki, T. Lefebure, B. Sket, and V.
706 Zaksek, V., 2009. A molecular test for cryptic diversity in ground water: how large
707 are the ranges of macro-stygobionts? *Freshwater Biology* 54:727-744.

708 Ward, J.V., and M.A. Palmer. 1994. Distribution patterns of interstitial freshwater
709 meiofauna over a range of spatial scales, with emphasis on alluvial river-aquifer
710 systems. *Hydrobiologia*. 287:147-156.

711

712

713

714 FIGURE LEGENDS

715 Figure 1: Distribution of groundwater fauna sampling stations and hydro-units within
716 the study region in the south-west of England.

717 Figure 2: Stygobitic species presence and community composition at sites within the
718 study region in the south-west of England. The size of the circles reflects the
719 number of species recorded.

720 Fig. 3: CCA ordination: Ordination by canonical correspondence analysis of
721 stygobiont species and stygophile data with significant environmental variables
722 (identified using the forward selection procedure) from boreholes, wells and springs
723 in SW England (Sept 2009-Nov 2011).

724

725 Table 1: Classification and description of the 5 hydro-units in the study area

Hydro-unit (no. of sites)	Geology	Main geologies	% Area coverage	Flow	Permeability (relative)
1 (10)	Granular	Quaternary & Palaeogene unconsolidated material	6.3	intergranular	Low-high
2 (37)	Igneous & metamorphic	Granite, basalt, spillite & gneiss & mica schists	8.6	Fracture	Low-mid
3 (57)	Mudstones & siltstones	Carboniferous, Triassic & Jurassic mudstones	51.6	Fracture and intergranular	Very low
4 (55)	Sandstones	Permo-Triassic, Jurassic & Devonian sandstones	17.2	Fracture and intergranular	Mid-high
5 (63)	Carbonate rocks	Chalk, Carboniferous & Devonian limestones	16.4	Karstic (solutional fissures and conduits)	Low-high

726

727

728 Table 2: Means and standard deviations of water chemistry parameters for each hydro-
 729 unit. Kruskal-wallis results indicate where determinands were significantly different
 730 between hydro-units.

731

Hydro-Unit	Statistic	EC μScm ⁻¹	DO mg/l	Temp °C	pH	DOC mg/l	NH ₄ mg/l	Cl mg/l	NO ₃ mg/l	SO ₄ mg/l	Ca mg/l	Mg mg/l	K mg/l	Na mg/l
1	n	8	8	8	8	8	8	8	8	8	8	8	8	8
	Mean	370.25	3.65	13.71	7.04	3.63	0.52	9.59	7.46	10.29	47.51	3.53	3.76	16.00
	Std Dev.	199.84	3.58	0.55	0.73	1.19	0.74	3.90	4.67	10.43	26.88	3.12	3.51	5.57
2	n	37	14	23	36	32	13	37	30	32	32	32	27	33
	Mean	276.14	8.91	9.65	6.35	1.41	0.02	8.86	4.66	4.42	25.07	12.33	4.95	20.46
	Std Dev.	193.32	3.41	1.36	0.76	1.72	0.01	7.18	4.85	3.36	12.37	13.98	6.11	14.47
3	n	47	29	29	45	47	29	46	41	47	46	46	45	45
	Mean	349.70	5.53	10.38	6.56	3.65	0.07	11.33	4.10	8.69	30.13	13.68	5.14	23.43
	Std Dev.	198.71	4.32	1.54	0.51	9.38	0.18	6.01	5.99	8.52	16.80	9.93	6.49	13.73
4	n	37	24	26	37	39	6	40	26	37	38	38	34	38
	Mean	423.89	5.90	10.60	6.81	11.52	0.82	12.14	9.07	11.04	37.68	7.40	5.92	18.22
	Std Dev.	245.74	3.89	1.44	0.75	15.48	1.63	6.77	8.30	12.87	27.10	5.21	7.63	9.23
5	n	54	25	39	54	54	17	55	53	52	53	52	43	53
	Mean	582.96	5.69	11.50	7.09	12.37	0.10	11.81	9.35	6.76	55.64	4.51	4.08	15.79
	Std Dev.	133.64	3.14	0.82	0.43	31.16	0.11	6.02	5.52	4.49	21.89	4.25	3.21	8.82
Kruskal Wallis Test	H Df p	48.54 4 <0.001	11.36 4 <0.001	22.37 4 <0.001	30.49 4 <0.001	53.51 4 <0.001	N/A	14.25 4 <0.01	38.66 4 <0.001	18.56 4 <0.01	44.73 4 <0.001	36.24 4 <0.001	5.51 4 0.239	13.44 4 <0.01

732 N/A – insufficient data (majority of values below detection limit).

733

734

735 Table 3: Abundance and composition of groundwater taxa (stygbionts and
 736 stygophiles) in boreholes (BH), wells and springs from the five hydro-units.
 737

Attributes of sample sites	Total	Sites sampled (no.)		
		BHs	Wells	Springs
No. of sites sampled:	221	124	56	41
<u>No. of sites in each hydro-unit (Hu)</u>				
Granular (Hu1)	10	9	1	0
Igneous/metamorphic (Hu2)	37	12	12	13
Mudstone/siltstone (Hu3)	56	28	19	9
Sandstone (Hu4)	55	34	11	10
Carbonate (Hu5)	63	41	13	9
No. of sites with GW fauna	155	70	51	34
Sites with no GW fauna* ¹	66	54	5	7
No. of sites with stygbionts	75	34	23	18

Taxa groups & species	Total Abundance	Percentage occurrence from all sites				Hydro-units present
Stygbionts	465	33.8	27.4	40.4	43.9	All
<u>Crustacea, Malacostraca, Eumalacostraca</u>						
<u>Peracarida</u>						
<u>Amphipoda</u>						
<u>Niphargidae</u>						
<i>Niphargus kochianus</i>	181	10.8	12.9	12.3	2.5	3, 5
<i>Niphargus fontanus</i>	28	5.4	4.8	7.0	4.9	1, 5
<i>Niphargus aquilex</i>	96	10.4	8.1	12.3	14.6	1, 2, 3, 4, 5
<i>Niphargus glenniei</i>	79	8.6	2.4	12.3	22.0	2, 3, 4, 5
<i>Microniphargus leruthi</i>	10	2.7	2.4	3.5	2.5	3, 4, 5
<u>Crangonyctidae</u>						
<i>Crangonyx subterraneus</i>	60	5.9	6.5	3.5	7.3	4, 5
<u>Isopoda</u>						
<u>Asellidae</u>						
<i>Proasellus cavaticus</i>	10	2.3	4.0	0	0	5
<u>Syncarida</u>						
<u>Bathynellacea</u>						
<u>Bathynellidae</u>						

<i>Antrobathynella stammeri</i>	1	0.5	0.8	0	0	4
Stygophiles	5318	63.5	44.4	86.0	90.2	All
<u>Crustacea, Maxillopoda</u>						
Copepoda	4181	47.3	26.6	70.2	78.0	1, 2, 3, 4, 5
<u>Crustacea, Ostracoda</u>	745	23.4	8.1	38.6	48.8	1, 2, 3, 4, 5
<u>Arthropoda, Arachnida, Acari, Trombidiformes</u>						
Hydrachnidae	182	16.2	10.5	21.1	26.8	1, 2, 3, 4, 5
<u>Annelida, Clitellata</u>						
Oligochaeta* ²	210	10.4	12.9	12.3	-	2, 3, 4, 5

738 *¹ Other freshwater (epigean) or terrestrial taxa may have been recorded at these sites

739 *² Not included where recovered from spring sources, due to presence of epigean species

740

741 Table 4: Summary of odds ratios from generalized linear models selected to describe
742 relationships between presence/absence data, hydro-units (Hu) and water chemistry.
743 (The level of significance reported in each model is represented by: * 0.05, ** 0.01,
744 ***0.001).

Odds ratios calculated relative to:

Hydro-unit 5 (carbonate)

Sample type 1
(Boreholes)

Generalized linear model (simplest model selected based on comparison of all model permutations)	Determinand (value)	Granular material (Hu1)	Igneous / metamorphic (Hu2)	Mudstone / siltstone (Hu3)	Sandstone (Hu4)	Sample type 2 (wells)	Sample type 3 (springs)
Pres/Abs~Hydro-unit	all data	0.84	0.80	0.12***	0.21***		
$\chi^2 = 33.8, df=5, p<0.001$							
	dissolved oxygen	2.00	1.00	0.16**	0.26*		
	nitrate	N/A	0.67	0.12***	0.28*		
	temperature	2.38	1.48	0.15**	0.23*		
Pres/Abs~Hydro-unit + value	calcium	0.97	0.98	0.18**	0.27**		
	chloride	0.93	0.63	0.11**	0.20**		
	pH	0.96	0.56	0.08***	0.16***		
Pres/Abs~Hydro-unit + sample type	Dissolved organic carbon	1.12	0.65	0.10***	0.21***	2.34*	3.01*
	elec. conductivity	1.09	0.59	0.10***	0.22**	2.19	3.07*
	magnesium	0.94	0.43	0.10**	0.19***	1.83	3.22*
	potassium	0.91	0.30*	0.07***	0.14***	1.77	4.52*
	sodium	1.00	0.49	0.12***	0.21**	1.92	3.20*
	sulphate	1.89	0.49	0.09***	0.19**	1.86	3.51*

746 N/A – excluded from analysis because insufficient reliable nitrate data available

747 Level of significance in model * 0.05, ** 0.01, ***0.001

748

749

750

751

752

753

754

755

756 Table 5: Partialled CCA results showing percentage of variation in species data explained
757 by environmental variables; hydro-units & groundwater chemistry, together and in isolation.

758

CCA	Constraint on Species data	Variable partialled out (covariable)	Sum of all canonical eigen-values (Ev)	Percentage variation (Ev x 100/4.998)	Fraction of variation explained by
CCA1	Chemistry	none	0.429	8.58%	-
CCA2	Hydro-units	none	0.628	12.57%	-
CCA3	Hydro-units	Chemistry	0.251	5.02%	Chemistry
CCA4	Chemistry	Hydro-units	0.376	7.52%	Hydro-units

Sum of all eigenvalues in a species only CA	4.998	
Overall amount of variation due to environmental variables	16.11%	CCA1 + CC4
Variation due to hydro-units & water chemistry together	3.56%	CCA1 –CCA3
Variation unexplained by hydro-units &/or water chemistry	83.89%	100 – (CCA1 + CCA4)

759

Figure 1: Groundwater faunal records and distribution of hydro-units within the study region in the south-west of England.





Figure 2: Stygobitic species presence and community composition at sites within the study region in the south-west of England. The size of the circles reflects the number of species recorded.



1 Running head: Groundwater faunal distributions

2

3 REGIONAL SCALE DRIVERS OF GROUNDWATER FAUNAL DISTRIBUTIONS

4

5 Tim Johns^{1,6}, J. Iwan Jones², Lee Knight³, Louise Maurice⁴, Paul Wood⁵, Anne
6 Robertson^{6*}

7 1. Environment Agency, Red Kite House, Howbery Park, Wallingford, Oxfordshire,
8 OX10 8BD, UK.

9 2. School of Biological and Chemical Sciences, Queen Mary University of London,
10 London E14NS

11 3. No. 1 The Linhay, North Kenwood Farm, Oxtou, Nr Kenton, Devon, EX6 8EX

12 4. British Geological Survey, Maclean Building, Crowmarsh Gifford, Wallingford,
13 Oxfordshire, OX10 8BB UK.

14 5. Centre for Hydrological and Ecosystems Science, Department of Geography,
15 Loughborough University, Loughborough, Leicestershire, LE11 3TU, UK

16 6* Department of Life Sciences, University of Roehampton, Holybourne Avenue,
17 London SW15 4JD, UK. T: +44 (0) 208 392 3456 E: a.robertson@roehampton.ac.uk

18 .*Author to whom correspondence should be addressed

19

20

21

22

23 *ABSTRACT*

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

Freshwater aquifers are a major source of drinking water; they also possess unique assemblages of organisms. However, little is known about the distributional drivers of obligate groundwater organisms at the regional scale. We examine the distribution and composition of stygobiont assemblages in a complex geological setting and explore the relationship between groundwater fauna, hydrogeology and water chemistry. In the study area we grouped similar geologies into five hydrogeological formations (hydro-units) within which habitats for groundwater fauna were broadly similar. We found that the occurrence of stygobionts differed significantly between hydro-units. Stygobionts were significantly less likely to be recorded in mudstone/siltstone and sandstone aquifers compared with carbonate rocks or with igneous/metamorphic rocks. Variance partitioning indicated that the hydro-units explained a greater proportion of the variance (7.52%) in the groundwater community than water chemistry (5.02%). However, much of the variation remained unexplained. The macrofaunal stygobiont species in our study area formed three groups: (1) *Niphargus glenniei* was recorded in a range of hydro-units but only in the west of the study area. (2) *Niphargus kochianus*, *Niphargus fontanus*, *Proasellus cavaticus* and *Crangonyx subterraneus* were predominately recorded in carbonate aquifers in the east of the study area. (3) *Niphargus aquilex* and *Microniphargus leruthi*, were found throughout the study area and in a range of hydro-units. We hypothesise that physical barriers exist that prevent some stygobiont taxa from colonizing apparently suitable geologies; the low permeability deposits dividing the western and eastern parts of the study area may partly explain the observed distributions.

49

50

51 *KEYWORDS*

52 Hydrogeology, Geology, British Isles, Devon, Dorset, Stygobionts, Stygophiles,

53 *Niphargus*, Water chemistry, groundwater.

54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78

INTRODUCTION

Freshwater aquifers represent approximately 65% of drinking water sources in the EU (Scheidleder and Visser, 2012), and thus their hydrogeology and chemistry are generally well known. In addition to their value as a resource, groundwaters possess unique assemblages of organisms and globally harbour a large reservoir of biodiversity with a high proportion of endemic and rare species compared with surface water habitats (Sket 2004, Gibert et al. 2009). In response to the energy poor nature of subterranean environments, groundwater invertebrates typically have low population densities and extended life histories with delayed maturity, greater longevity, smaller egg-clutch size, larger eggs and lower percentages of mature ovigerous females than related surface-dwelling (epigean) species (Gibert et al. 1994). The low population numbers and slow reproduction of groundwater organisms may result in an increased vulnerability to anthropogenic impacts compared to epigean taxa. This vulnerability may be compounded by their slow dispersal rates (Culver et al. 2009).

Although research centred on surface-groundwater interactions is burgeoning and has moved from descriptive studies to a more experimental approach (introduction to this special issue, Larned 2012), research into deeper groundwaters is less developed, perhaps because of the difficulties in accessing this environment. Despite numerous studies describing the occurrence and distribution of groundwater fauna (e.g. Botosaneanu 1986, Proudlove et al. 2003, Robertson et al 2009, Gibert et al. 2009), relatively little is known about the environmental and/or historical drivers of their distribution patterns. The relative importance of a given driver is likely to vary across spatial and temporal scales. At the continental scale, stygobiont distribution

79 patterns have been shaped by events in deep geological time, for example the
80 movement of continental plates (e.g. Schminke 1974), orogenic and eustatic events
81 (Stock 1980, Boutin and Coineau 1990, Notenboom 1991), and the alternation of
82 marine transgressions and regressions (e.g. Jaume and Humphreys 2001). Long
83 term climate change has also shaped the distribution of groundwater taxa. In the
84 northern hemisphere the Pleistocene glaciations appear to have had a catastrophic
85 effect on groundwater communities, leaving Northern Europe with a depauperate
86 groundwater fauna compared to the rest of Europe. For example, only eight species
87 of obligate groundwater macro Crustacea have been recorded from mainland Britain,
88 including four species of *Niphargus*, the most speciose groundwater genus
89 (Proudlove et al. 2003, Robertson et al. 2009, Knight and Gledhill 2010). This
90 contrasts with 134 species of *Niphargus* recorded in the Balkans biogeographical
91 area (McInerney et al. 2014). In Northern Europe, contemporary distributions reflect
92 patterns of past glacial activity (e.g. Strayer 1994, Robertson et al. 2009). However,
93 there are some species present in formerly deglaciated areas, and those species
94 that remain in Northern Europe must have survived the impacts of glacial and
95 periglacial conditions (Holsinger 1988, Proudlove et al. 2003, Kristjansson and
96 Svavarsson 2007, Lefebure et al. 2007, McInerney et al. 2014). Indeed, recent
97 research has shown that the small number of *Niphargus* species that are present in
98 the UK are ancient endemics that have survived millions of years of extreme climate
99 changes (McInerney et al. 2014). At smaller scales, stygobiont distribution patterns
100 are influenced by the availability of suitable habitat and aquifer connectivity as well
101 as the ecological tolerances, competitive interactions and dispersal abilities of
102 species (Gibert et al. 1994, Castellarini et al. 2007, Hahn and Fuchs 2009).

104 Regional investigations that have gathered large datasets on stygobiont presence
105 and environmental conditions are relatively rare (Mauclaire and Gibert 2001, Galassi
106 et al. 2009, Martin et al. 2009, Dole-Olivier et al. 2009b, Hahn and Fuchs 2009). In
107 these studies, geology appears to be an important determinant of stygobiont
108 distributions because it controls the availability of suitable habitats (Datry et al. 2005,
109 Dole-Olivier et al. 2009b, Botosaneanu 1986, Hahn and Fuchs 2009). Most
110 stygobionts are dependent on well-developed void spaces with wide pore throats or
111 fracture apertures. These features commonly occur in karstic, porous and fractured
112 aquifers and this is where obligate groundwater communities are most diverse and
113 abundant (e.g. Gibert and Deharveng 2002, Dole-Olivier et al. 2009b). In addition to
114 providing physical habitat, the degree to which void spaces are developed and
115 connected also influences the exchange of oxygen, nutrients and organic matter with
116 the surface, and thus the availability of food for the groundwater community.

117 Observed distribution patterns may also relate to the opportunities for dispersal. In
118 groundwaters the habitat is very fragmented, limiting the dispersal of groundwater
119 fauna (Lefebure et al. 2007). Stygobiont assemblages are therefore characterised
120 by endemic species with limited geographical ranges. Even species which are
121 apparently widely distributed have emerged as cryptic lineages (Lefebure et al. 2007,
122 Trontelj et al. 2009, McInerney et al. 2014). Habitat fragmentation in groundwaters
123 has been most frequently discussed in relation to karst habitats (Trontelj et al. 2009,
124 Modovan et al. 2012), however it may also occur as a result of the juxtaposition of
125 different geologies. The presence of mudstones with low permeability and restricted
126 void spaces may form a physical barrier to stygobiont dispersal (Hahn and Fuchs
127 2009). Some stygobiont taxa, however, may be able to exploit superficial deposits or
128 the hyporheic zones of rivers as a means of dispersal (e.g. Ward and Palmer 1994,

129 Castellarini et al. 2007). The water chemistry of an aquifer, e.g. concentrations of
130 dissolved organic carbon and nitrates, may also correlate with the distribution of
131 groundwater taxa (Martin et al. 2009).

132

133 In this study we address the limited understanding of the drivers of groundwater
134 faunal distributions at the regional scale by carrying out an extensive sampling
135 programme within the complex geological setting of the south-west of the UK. Our
136 aims were to determine the detailed spatial distributions of the UK stygobiont species
137 in the study area and investigate whether the distribution and composition of
138 stygobiont assemblages varies with hydrogeology and water chemistry. We build on
139 the design of the PASCALIS project on European groundwater biodiversity which
140 adopted a hierarchical sampling strategy and included two types of aquifer (porous
141 and karstic) (Dole-Olivier et al. 2009a). Our approach was to group the different
142 hydrogeological formations present in the study area into five 'hydro-units' where the
143 different geologies within a given hydro-unit provide broadly similar habitats for
144 groundwater fauna. We sampled groundwater assemblages within the different
145 hydro-units to determine whether species were restricted to particular hydro-units,
146 and sampled water chemistry to determine whether faunal distributions reflect local
147 variation in water chemistry.

148

149

METHODS

Study area and hydrogeological units

151 The study area in south west England is an area of contrasting topography and
152 geology dominated by lowland (max. elevation 279 m) sedimentary rocks in the east
153 and higher elevation (max. 621 m) metamorphic and igneous rocks in the west (Fig.

154 1). The region's annual average rainfall is approximately 900-1000 mm
155 (<http://www.metoffice.gov.uk/climate/uk/sw/>).

156

157 Within the study area 271 different geological formations have been mapped at the
158 1:50,000 scale (Reproduced from the British Geological Survey Map data at the
159 original scale of 1:50 000 Licence 2011/057 British Geological Survey. ©NERC. All
160 rights reserved). To assess the general hydrogeological controls on species
161 distributions, these geologies were grouped into categories termed "hydro-units".
162 The hydro-units are based on broad differences in geology and groundwater flow
163 characteristics which result in different groundwater habitats for fauna. There are
164 two main types of groundwater flow: intergranular and flow through fractures (Price
165 1996). Unconsolidated granular deposits in which all groundwater movement is
166 intergranular were grouped together as hydro-unit 1 because these deposits form a
167 habitat in which fauna can only live within the pore spaces between the sediment
168 grains. Fractured rocks may be igneous, metamorphic or sedimentary in origin.
169 Igneous and metamorphic rocks (which are of volcanic origin) were grouped together
170 (hydro-unit 2) because there were insufficient data to determine differences in habitat
171 type between them. They were grouped separately from other strata because
172 groundwater flow is only through fractures, and due to of their volcanic origin, they
173 have a bedrock geochemistry that is different to other consolidated rocks.
174 Sedimentary rocks may be non-carbonate or carbonate. Non-carbonate
175 sedimentary rocks often have varying degrees of consolidation and, therefore, form
176 aquifers in which flow may be both inter-granular in the less consolidated areas and
177 through fractures in consolidated rock. They are classified according to grain size as
178 mudstones, siltstones and sandstones (Price 1996). Sandstones form aquifers with

179 much higher permeability than siltstones and mudstones which often form aquitards
180 or aquicludes (Price 1996, Allen et al. 1997). The difference in permeability is likely
181 to result in both a different groundwater chemistry and a different physical habitat for
182 fauna. They were therefore separated into a mudstone/siltstone hydro-unit (3) and a
183 sandstone hydro-unit (4). Hydro-unit 5 includes all carbonate geologies. In these
184 rocks, fractures are enlarged by dissolution to form fissures and caves providing
185 substantially larger and more inter-connected void spaces than any other rock type,
186 which can serve as habitats for fauna. In the study area carbonate rocks range from
187 the weakly karstified Cretaceous Chalk (Maurice et al, 2006) to the highly karstified
188 limestones of Devonian and Carboniferous. Where the Chalk overlies the Upper
189 Greensand Formation these strata are commonly in hydraulic continuity (Allen et al.
190 1997). This means that it is likely that fauna may be living in both the Chalk and the
191 Upper Greensand, and that it is not possible to distinguish the two habitats. The
192 Upper Greensand Formation is a non-karstic aquifer comprising sandstones and
193 sands, but where it is in hydraulic continuity with the Chalk it was included in the
194 carbonate hydro-unit (5). Elsewhere, outcrops of the Upper Greensand Formation
195 are isolated from the Chalk and, were therefore grouped with the other sandstones in
196 the sandstone hydro-unit (4). The properties of the five hydro-units are summarized
197 in Table 1.

198

199 The distribution and extent of the different hydro-units varied across the study area
200 (Fig. 1). In particular there was a pronounced difference between the hydro-units in
201 the east and west of the study area. Mudstones/siltstones (hydro-unit 3), covered the
202 largest surface area in the study region and were most common in the west.
203 Sandstones (hydro-unit 4), were the next most common, and occurred across the

204 region, with the largest outcrops in the west and north-west. Carbonate aquifers
205 (hydro-unit 5), were predominately located in the east, where there were extensive
206 Cretaceous chalk downlands. In the west carbonate rocks were limited to much
207 smaller isolated outcrops of older limestones, separated by mudstones and
208 sandstones. Granular material (hydro-unit 1) was restricted to the far south eastern
209 edge of the study area, whereas igneous and metamorphic rocks (hydro-unit 2),
210 were only present in the west and south west.

211

212 *Sampling*

213 In total, 221 sites were sampled. The number of samples taken within each hydro-
214 unit was approximately proportionate to the surface area it occupied (Table 1). Within
215 each unit, sites were selected to maximize spatial coverage. Boreholes free from fine
216 mesh well screens and in-situ equipment were preferentially selected. Borehole and
217 well sampling sites ranged from 0.36 to 130 m deep and 0.05 to 2 m in diameter.
218 Where there were insufficient boreholes or wells, springs were sampled. Each site
219 was sampled on a single occasion. Most samples (197) were collected between
220 September 2009 and May 2010, and water chemistry measurements were made at
221 the time of sampling. A further 24 samples were collected in November 2011; no
222 water chemistry data were obtained for these samples.

223

224 Fauna were collected from boreholes and wells using a weighted plankton net (63µm
225 mesh, diameter varied from 35 – 300mm dependent on borehole aperture). The net
226 was lowered to the base of the borehole/well and moved up and down vigorously to
227 disturb the sediment before retrieval. For springs, a net was swept through the
228 sediment of the spring chamber or at the spring head source. The samples were

229 preserved in >90% ethanol and all equipment thoroughly cleaned between sites. In
230 the laboratory animals were separated from debris, stygobionts were identified to
231 species level using Gledhill et al. (1993) and Knight and Gledhill (2010); Ostracoda
232 (identified to class), Hydracarina (identified to clade) and Copepoda and Oligochaeta
233 (identified to sub-class) were probably predominantly stygophiles and stygoxenes.
234 Freshwater epigeal taxa and terrestrial taxa were noted.
235 Electrical conductivity, pH, dissolved oxygen and temperature were recorded using a
236 multi-parameter probe (YSI 600QS) lowered to the base of each borehole/well. The
237 probe was calibrated prior to each sampling trip. Water samples for laboratory
238 analysis were taken using a bailer. At springs, field chemistry and water samples
239 were taken directly from the spring head or catchpit. Water samples were filtered and
240 refrigerated before dispatch to analytical laboratories within 48 hours. Analysis for
241 major anions (Cl^- , SO_4^{2-} , NO_3^-) and cations (Mg^{2+} , K^+ , Na^+ , Ca^{2+}) was performed
242 using high pressure liquid chromatography using a Dionex ED40 electrochemical
243 detector (min. reporting value; 1.0mg l^{-1}). Ammoniacal nitrogen was analyzed by
244 colourimetric analysis using a Konelab discrete analyser (min. reporting value; 0.03
245 mg l^{-1}). Dissolved organic carbon (DOC) was analyzed by a Skalar San Plus 1050 air
246 segmented flow analyser (min. reporting value 0.2 mg l^{-1}).

247

248 *Data analysis*

249 Chemical data were tested for normality (Q-Q plots) and homogeneity of variance
250 (Fligner-Killeen test), and a non-parametric ANOVA was used to investigate
251 differences in water chemistry between the five different hydro-units (selected as
252 factors). A generalized linear model with a binomial error structure was then used to
253 examine the relationships between presence/absence of stygobiont taxa, hydro-units

254 and water chemistry. Statistical analyses were performed using R version 3.0.2
255 (2013), using standard packages plus the package for analysis of over dispersed
256 data (Lesnoff et al, 2012). The initial model selected hydro-unit as a factor. The
257 model was then run with all stygobiont presence/absence data, using the carbonate
258 hydro-unit (5) as the point of comparison because it: i) comprised the most sites, ii)
259 was the strata in which stygobionts were most common, and iii) also yielded a broad
260 range of water quality data. Models were then run for each set of water quality data
261 (i.e. each chemical determinand with associated presence/absence and hydro-unit
262 data), the odds ratios were calculated and predicted probabilities of stygobiont
263 presence were determined from the mean value for each determinand. Different
264 permutations of the model were run to explore whether the inclusion of sample type
265 (borehole, well or spring) and/or water quality data improved the model, and to
266 identify any interactions between factors. Outputs were checked to verify the data
267 were not over dispersed and models were then compared, using a one-way ANOVA,
268 to identify any significant differences between models. Where there was no
269 difference, the simplest model was selected and the odds ratios calculated to
270 describe the relative likelihood of recording stygobionts in each hydro-unit.

271

272 The association between the faunal community and environmental variables was
273 investigated using Canonical Correspondence Analysis (CCA) within Canoco for
274 Windows Ver.4.51 (Ter Braak and Smilauer 1998). Only those sites with concurrent
275 chemistry analysis (190 samples) were used in this analysis and sites where there
276 was uncertainty over the geological unit from which the sample was obtained (7
277 samples) were excluded.

278 A series of ordinations were undertaken including groundwater faunal abundance,
279 hydro-units and water chemistry data. Both stygobiont and stygophile taxa were
280 included in the analysis to reflect the diversity and distribution of the community
281 recorded from each sample location where fauna were recorded, although the
282 results and discussion are focused on the stygobiont taxa. To determine how
283 individual variables might explain patterns in the faunal data (stygobionts and
284 stygophiles), we applied the forward selection procedure ($P \leq 0.05$ after Bonferonni
285 correction). Only those variables having a significant association with the faunal
286 community distribution were included in the final analysis and presented in the output
287 diagrams. The stygobiont *Antrobathynella stammeri* (Jakobi, 1954), which was
288 recorded only once, was fitted passively to the ordination. To determine the
289 proportion of variance in the faunal data explained by the hydro-units and water
290 chemistry data, a variance partitioning approach was used (Borcard 1992). This
291 constrained the species ordination for each of these co-variable groups in turn, while
292 partialling out the other variables from the ordination.

293

294

295

RESULTS

296 As expected water chemistry varied significantly between the five hydro-units,
297 reflecting differences in bedrock geochemistry and permeability. With the exception
298 of potassium, significant differences were recorded between each chemical
299 determinand and the five hydro-units (Table 2).

300

301 Stygobionts and/or stygophiles were recorded from 70% of the study sites (Table 3)
302 and all hydro-units (Fig.1). The remaining 30% of sites contained no groundwater

303 taxa although epigean taxa were recorded at some sites. Stygophiles were more
304 abundant and recorded more frequently than stygobionts (59% compared to 34% of
305 sites). All of the stygobiont macro-Crustacea known to occur in the UK were
306 recorded, although their presence and community composition varied considerably
307 across the study region. Boreholes in the carbonate hydro-unit (unit 5) supported the
308 most diverse stygobiont communities, with up to four taxa recorded in a single
309 borehole, compared to a maximum of two taxa from other hydro-units (Fig. 2).

310

311 *Niphargus aquilex* (Schiödte, 1855), was recorded throughout the study region and
312 from all hydro-units (Fig. 2). In contrast, *Niphargus glenniei* (Spooner, 1952) was
313 only recorded in the west of the study area despite being recorded in most hydro-
314 units. *Niphargus kochianus* (Bate, 1859), *Niphargus fontanus* (Bate, 1859) and
315 *Proasellus cavaticus* (Leydig, 1871) were primarily recorded in the carbonate hydro-
316 unit (5) in the east of the region (Fig. 2), but were absent from this hydro-unit and all
317 other hydro-units in the west of the study area. *Crangonyx subterraneus* (Bate,
318 1859) and *Microniphargus leruthi* (Schellenberg 1934) were rare, and were primarily
319 recorded in the east (Fig. 2). *Antrobathynella stammeri* (Jakobi, 1954) was recorded
320 once.

321

322 The occurrence of stygobionts differed significantly between hydro-units [$\chi^2=33.8$,
323 $df=5$, $p<0.001$, Table 4]. From the odds ratio (OR), stygobionts were 88% less likely
324 to be recorded in mudstone/siltstones (hydro-unit 3) [OR=0.12, 97.5%CI 0.034-
325 0.030, $p<0.001$] than in carbonates (hydro-unit 5) and about 80% less likely to be
326 recorded in sandstones (hydro-unit 4) [OR=0.20, 97.5%CI 0.08-0.50, $p<0.001$], than
327 carbonates (hydro-unit 5). Stygobionts were also less likely to be recorded in

328 mudstones/siltstones (hydro-unit 3) and sandstones (hydro-unit 4) than in
329 igneous/metamorphic sites (hydro-unit 2) and granular sites (hydro-unit 1). All
330 differences were significant except for sandstone sites relative to granular sites.
331 The generalized linear model generally improved (based on the AIC scores and level
332 of significance) with the addition of either sample type or a water chemistry
333 determinand as a value. Mudstones/siltstones and sandstones (hydro-units 3 and 4)
334 were significantly different from carbonate rocks (hydro-unit 5) throughout the
335 iterations. No significant interactions between factors (e.g. hydro-unit combined with
336 sample type) were identified. By including water chemistry determinands with
337 presence/absence in the model, improved models were obtained with calcium,
338 chloride and pH , indicating a potential association of these variables with stygobiont
339 presence. The predicted probability of stygobiont presence in any hydro-unit was
340 higher with increasing calcium concentrations. In contrast, the predicted probability
341 of stygobiont presence in any hydro-unit declined as chloride concentrations
342 increased. There was no clear relationship between pH and the presence of
343 stygobionts.

344

345 The inclusion of sample type significantly improved the models that included
346 electrical conductivity, magnesium, potassium, sodium, sulphate and dissolved
347 organic carbon. For the model with potassium, the igneous/metamorphic aquifers
348 (hydro-unit 2) were identified as significantly different when sample type was
349 included. The odds ratio of recording stygobionts in this unit was also lower
350 compared with the simplest model where no additional factors were included. Overall
351 there was a greater likelihood of recording stygobionts in springs (sample type 3)
352 than boreholes (sample type 1: Table 4). However, presence or absence of

353 stygobionts across the hydro-units was significant ($p < 0.001$) irrespective of the
354 inclusion of sample type in the model.
355
356 Canonical correspondence analysis indicated a separation of the sample centroids
357 for the five hydro-units based on the faunal community data (Fig. 3), the first two
358 canonical axes explained 20.0% and 13.6% of the total variance in faunal
359 abundance respectively (Fig. 3). These two axes combined explained 78.4% of the
360 variance in the species-environment relationship (46.7% and 31.7% respectively).
361 Water temperature ($^{\circ}\text{C}$), magnesium (mg/l) and potassium (mg/l) were significantly
362 correlated with the distribution of the groundwater fauna (all $p < 0.01$). The stygobionts
363 *Niphargus kochianus*, *Microniphargus leruthi*, *Crangonyx subterraneus* and
364 *Proasellus cavaticus* were associated with the centroids for the sandstone and
365 carbonate hydro-units (hydro-unit 4 and 5). In contrast, some taxa were not strongly
366 associated with the centroids of any single hydro-unit but were associated with
367 higher concentrations of potassium (*N. fontanus* and *N. aquilex*) and higher water
368 temperature (*N. glenniei*). Variance partitioning indicated that 7.5% of the variance in
369 faunal abundance could be explained by the hydro-units and 5.0% by water
370 chemistry (Table 5), whilst 3.6% of the variance was shared by hydro-units and water
371 chemistry together.

372

373

DISCUSSION

374 Stygobionts and stygophiles were found throughout the region and were present in
375 all hydro-units. However, stygobionts were absent from 70% of sites. Eberhard et al.
376 (2009) and Hancock and Boulton (2009) found that repeated sampling over time at
377 groundwater sites increased the number of species recorded, thus we cannot

378 exclude the possibility that the low frequency of stygobiont occurrence in our study
379 may be an artifact of low sampling effort, as we sampled each site on only one
380 occasion.

381 We found that stygobiont occurrence and community composition differed between
382 hydro-units. Stygobionts were significantly more likely to be recorded in the
383 carbonate and igneous/metamorphic hydro-units, than in the mudstone/siltstone or
384 sandstone units. Several stygobiont species (*Niphargus kochianus*, *Microniphargus*
385 *leruthi*, *Crangonyx subterraneus* and *Proasellus cavaticus*) were predominantly
386 associated with the carbonate hydro-unit. Other stygobionts (*Niphargus aquilex* and
387 *Niphargus glenniei*) were found across a range of hydro-units.

388 The carbonate hydro-unit also supported the most diverse stygobiont communities,
389 with up to four stygobiont species co-occurring in a single borehole. Carbonate
390 aquifers are characterised by solutional fissures and conduits providing connected
391 water-filled void spaces which are sufficiently large for groundwater organisms to
392 inhabit (Humphreys 2009). Furthermore, their highly permeable nature may result in
393 a greater supply of food from the surface, permitting the co-existence of more
394 species than in less permeable hydro-units, where food shortages may be severe
395 (Malard et al. 2009). Studies in continental Europe have demonstrated previously
396 that karst aquifers support some of the most diverse stygobiont communities (Dole-
397 Olivier et al., 2009b; Martin et al., 2009; Galassi et al., 2009; Hahn and Fuchs, 2009).

398 The sandstone hydro-unit was also anticipated to provide a suitable habitat for
399 diverse stygobiont communities because this has been the case in other regions
400 (Humphreys 2009). Therefore, the low numbers of individuals and low diversity
401 recorded in our study was surprising. One explanation is that the Permo-Triassic

402 sandstones, which comprise part of this hydro-unit contain lower permeability
403 mudstone beds in this area (Allen et al. 1997). This may result in a fragmented
404 habitat with areas containing poorly developed fracture networks providing a poor
405 habitat. The deposits are also affected by faulting which may result in further
406 barriers to stygobiont movement. However, the sandstone (hydro-unit 4) is
407 predominantly distributed in the west of the study area, which is outside the
408 observed range of *Niphargus kochianus*, *Niphargus fontanus* and *Proasellus*
409 *cavaticus* (Fig. 2), three of the most common stygobite species in the east of the
410 study area. It is not clear whether the absence of these species in the west is
411 because the sandstones do not form a suitable habitat for them, or because they
412 have not dispersed to these areas.

413 The mudstone/siltstone hydro-unit had the lowest occurrence of stygobionts
414 suggesting that it provides a less suitable habitat, perhaps as a result of the small
415 and often weakly connected voids, and the low levels of oxygen and nutrients. Our
416 findings are consistent with those of Hahn and Fuchs (2009), who also reported that
417 mudstones were characterized by depauperate communities.

418

419 Our results demonstrate that the five hydro-units had different water chemistries.
420 Therefore it was not possible to determine whether differences in groundwater fauna
421 occurred due to water chemistry differences between the units, or differences
422 between the physical voids present within the different units (Maurice and
423 Bloomfield, 2012). However, in this study, groundwater taxa occurred in all units and
424 the proportion of the variance in the abundance of the groundwater community
425 explained by water chemistry was smaller than that of the hydro-units. The
426 mudstone/siltstone and sandstone units remained significantly different from the

427 carbonate unit in the general linear model, irrespective of the addition of water
428 chemistry parameters. However, the igneous/ metamorphic unit only became
429 significantly separated from the other hydro units when potassium (together with
430 sample type) was included in the model. The higher potassium concentrations
431 reported from the igneous/ metamorphic hydro-unit relative to the carbonate hydro-
432 unit reflects the contrasting water chemistry of these different geologies. These
433 findings provide evidence that suggests that, across our study sites, water chemistry
434 had less influence over the distribution of groundwater taxa than physical
435 hydrogeological characteristics. Other studies have also found little relation between
436 environmental variables, water chemistry and the distribution of groundwater taxa at
437 a regional scale (e.g. Dumas and Lescher-Moutoue 2001, Paran et al. 2005, Dole-
438 Olivier et al. 2009b, Galassi et al. 2009). Nevertheless, in all these studies there are
439 considerable difficulties when trying to separate the influence of potentially
440 confounding factors that occur over a range of geographical scales that may
441 determine stygobiont distributions (Dole-Olivier et al., 2009b; Stoch and Galassi,
442 2010; Maurice and Bloomfield, 2012).

443

444 The probability of finding stygobionts appeared to increase with an increase in
445 calcium concentrations. This is probably because stygobionts were most often found
446 in the carbonate hydro-unit where calcium concentrations are high. Although
447 calcium is necessary for crustacean carapace development (Rukke, 2002), it does
448 not appear to be limiting in our study area as high stygobiont frequencies were also
449 recorded from the igneous/metamorphic hydro-unit, where the mean calcium
450 concentration was the lowest of all the hydro-units sampled.

451

452 In contrast to Dole-Olivier et al. (2009b), we found no significant relationship
453 between dissolved oxygen concentration and stygobiont occurrence. This is
454 probably because all but 14 of our sites had oxygen concentrations $>1 \text{ mg l}^{-1}$, which
455 is the threshold below which oxygen has been reported to be critical for groundwater
456 taxa (Malard and Hervant 1999, Hahn 2006). In addition, stygobionts are widely
457 acknowledged to be tolerant of low oxygen conditions in comparison to epigean
458 species (Malard and Hervant 1999).

459 The greater likelihood of finding stygobionts in spring sites (sample type 3) than in
460 borehole sites (sample type 1) may be because it is easier to capture fauna in
461 springs than in boreholes, and also because springs form a distinct ecotone where
462 there are higher nutrients and more diverse communities than in deeper
463 groundwaters (Smith et al. 2003) However, the use of springs as an explanatory
464 variable did not fundamentally modify or significantly influence the differences we
465 identified between the hydro-units.

466

467 Most of the variation in the occurrence of groundwater fauna across the study region
468 was not explained by either the hydro-units or water chemistry. One of our most
469 striking findings was the differing distribution patterns of stygobiont species within the
470 study area. Most stygobiont species had restricted, and different distributions. The
471 exception was *Niphargus aquilex* which was recorded from all hydro-units and
472 across the whole study area. This species is widely distributed in central and
473 southern Europe also (Botosaneanu 1986), with an apparent range that spans $>$
474 2300 km east – west (Trontelj et al. 2009). In the British Isles it has been extensively
475 recorded in riverine hyporheic zones and is particularly associated with shallow
476 alluvial aquifers (Proudlove et al. 2003) and, thus, may have dispersed across the

477 area using this route (the hyporheic corridor concept of Stanford and Ward 1993).
478 Nevertheless, recent research shows that the wide distribution of *Niphargus aquilex*
479 is an artifact generated by cryptic diversity; the complex contains seven cryptic taxa,
480 two of which are unique to the British Isles (McInerney et al. 2014). Other widely
481 distributed *Niphargus* “species” have also been found to be species complexes (e.g.
482 *N. virei*, Lefebure et al. 2006 and *N. rhenorhodanensis*, Lefebure et al 2007, see also
483 Trontelj et al. 2009).

484

485 Although there would appear to be suitable habitats distributed across the study
486 area, other stygobiont species exhibited distinct spatial segregation. The endemic
487 *Niphargus glenniei* was only recorded in the west and *Niphargus kochianus*,
488 *Niphargus fontanus*, and *Proasellus cavaticus* were only found in the east. One
489 possible explanation for these observed distributions is that they result from biotic
490 interactions between the species. *Niphargus glenniei* for example, did not co-occur
491 with either *Niphargus fontanus* or *Niphargus kochianus*. *Niphargus fontanus* is much
492 larger and may have a competitive or predatory advantage. However, stygobiont
493 species can co-exist, and up to four species were found in a single borehole in the
494 east of our study area. Although in our study *Niphargus glenniei* was mostly
495 recorded alone or occasionally together with the smaller species *Microniphargus*
496 *leruthi*, other studies have reported it with the larger species *Niphargus aquilex*
497 (<http://hcrs.freshwaterlife.org/>). An alternative hypothesis is that geological barriers
498 are present that constrain the dispersal of these organisms, as has been reported for
499 other groundwater taxa (Gooch and Hetrick 1979; Goricki and Trontelj 2006).
500 Trontelj et al. (2009) observed that many groundwater taxa appear to be able to
501 disperse and yet are confined by boundaries beyond which dispersal is impossible

502 (although the nature of these boundaries is currently unclear). The
503 mudstones/siltstones and other low permeability rocks across the centre of our study
504 area may form a barrier between suitable stygobiont habitats in the east and west,
505 and could explain the different species distributions. However, given the occurrence
506 of all stygobiont species in spring habitats, it is surprising that groundwater species
507 have not been able to utilize shallow superficial deposits (e.g. hyporheic zone
508 corridors) to disperse between the east and west of the study area.

509 The life histories and restricted distributions of groundwater taxa mean that they are
510 vulnerable to anthropogenic change (Gibert et al. 1994). *Niphargus glenniei*, which is
511 endemic to two counties in southern England, has been designated a UK
512 Biodiversity Action Plan species. However, other groundwater taxa have no such
513 recognition, and European groundwater monitoring programmes do not consider
514 groundwater ecology. Identifying the drivers of observed distribution patterns in
515 groundwater taxa is important because this information may then be used to inform
516 the development of management plans for the conservation of these unique
517 assemblages.

518 .

519 CONCLUSIONS

520 In our regional scale study, there were significant differences in the groundwater
521 assemblages present in geologies with different hydrogeological characteristics.
522 Stygobionts occurred most frequently and were most diverse in the carbonate hydro-
523 unit (5) which is characterized by karstic, solutionally enlarged voids providing an
524 extensive physical habitat, and a good supply of nutrients and oxygen. They were
525 also relatively abundant and diverse in the igneous and metamorphic hydro-unit (2)
526 and granular aquifers (hydro-unit 1) but were relatively rare in the sandstone hydro-

527 unit (4). They occurred least frequently in the mudstone/ siltstone hydro-unit (3)
528 which has low permeability resulting in poor physical and water chemistry conditions
529 for stygobionts. The hydro-units explained a greater proportion of the variation in
530 stygobiont data than water chemistry, suggesting that physical hydrogeology may be
531 more important than water chemistry in determining stygobiont distributions.
532 However, a large proportion of the variation in stygobiont abundance was not related
533 to either the hydro-units or water chemistry and we were intrigued to find that some
534 stygobiont species were spatially segregated within our study area. It appears that
535 barriers may exist that prevent some stygobiont taxa from colonizing apparently
536 suitable geologies, and the low permeability deposits dividing the western and
537 eastern parts of the study area may partly explain the observed distributions. The
538 stygobiont species fell into three contrasting groups: (1) *Niphargus glenniei* was only
539 found in the west of the study area, but was found in a range of hydro-units with
540 varying void types and water chemistry. (2) *Niphargus kochianus*, *Niphargus*
541 *fontanus*, *Crangonyx subterraneus* and *Proasellus cavaticus* were only found in the
542 east of the study area and predominantly in carbonate aquifers. (3) *Niphargus*
543 *aquilex* and *Microniphargus leruthi*, were found throughout the study area and in a
544 range of hydro-units suggesting that they have good dispersal abilities and may be
545 able to circumvent low permeability barriers, perhaps by utilizing superficial deposits
546 or hyporheic zone corridors. Identifying the precise nature of the barriers to dispersal
547 and the development of monitoring and conservation plans must be a priority for
548 these restricted and vulnerable species.

549

550

ACKNOWLEDGMENTS

551 This research was funded by the Esmee Fairbairn Foundation, grant number 08-
552 101,7 to ALR, PJW, LK and LM. We thank the many landowners who gave us
553 permission to cross their land, and the Environment Agency for their help and
554 assistance in accessing sites and provision of equipment. L. Maurice publishes with
555 the permission of the Executive Director, British Geological Survey (NERC).

556

557

LITERATURE CITED

- 558 Allen, D.J., L.J. Brewerton, L.M. Coleby, B.R. Gibbs, M.A. Lewis, A.M. MacDonald,
559 S.J. Wagstaff, and Williams, A.T., 1997. The physical properties of the major
560 aquifers in England and Wales. BGS Technical Report WD/97/34, Environment
561 Agency R&D Publication 8. 312 pp
- 562 Botosaneanu, L. 1986. *Stygofauna Mundi*. E.J. Brill, Leiden.
- 563 Borcard, D., P. Legendre, and P. Drapeau, 1992. Partialling out the spatial
564 component of ecological variation. *Ecology* 73:1045-1055.
- 565 Boutin, C., and N. Coineau. 1990. "Regression model, "modèle biphasé" d'évolution
566 et origine des microorganismes stygobies interstitiels continentaux. *Revue de*
567 *Micropaléontologie* 33:303–322
- 568 Castellarini, F., F. Malard, M-J. Dole-Olivier, and J. Gibert. 2007. Modelling the
569 distribution of stygobionts in the Jura Mountains (eastern France). Implications for
570 the protection of ground waters. *Diversity and Distributions* 13:213-224.
- 571 Culver D.C., T. Pipan, and K. Schneider. 2009. Vicariance, dispersal and scale in
572 the aquatic subterranean fauna of karst regions. *Freshwater Biology* 54:918-929.
- 573 Datry, T., F. Malard, and J. Gibert. 2005. Response of invertebrate assemblages to
574 increased groundwater recharge rates in a phreatic aquifer. *Journal of the North*
575 *American Benthological Society* 24:461-477.

576 Dole-Olivier, M.-J., F. Castellarini, N. Coineau, D.M.P. Galassi, P. Martin, N. Mori, A.
577 Valdecasas, and J.Gibert. 2009a. Towards an optimal sampling strategy to
578 assess groundwater biodiversity: comparison across six European regions.
579 *Freshwater Biology* 54:777-796.

580 Dole-Olivier, M.-J., F. Malard, D. Martin, T. Lefebure, and J. Gibert. 2009b.
581 Relationships between environmental variables and groundwater biodiversity at
582 the regional scale. *Freshwater Biology* 54:797-81

583 Dumas, P., and F. Lescher-Moutoue. 2001. Cyclopid distribution in an agriculturally
584 impacted alluvial aquifer. *Archiv für Hydrobiologie* 150:511-528.

585 Eberhard, S. M., S.A. Halse, M.R. Williams, M.D. Scanlon, J. Cocking, and H.J.
586 Barron. 2009. Exploring the relationship between sampling efficiency and short-
587 range endemism for groundwater fauna in the Pilbara region, Western Australia.
588 *Freshwater Biology* 54:885-901.

589 Galassi, D.M.P., F. Stoch, B. Fiasca, T. Di Lorenzo, and E. Gattone. 2009.
590 Groundwater biodiversity patterns in the Lessinian Massif of northern Italy.
591 *Freshwater Biology* 54:830-847.

592 Gibert, J., D. Danielopol, and J. Stanford (Eds). 1994. *Groundwater Ecology*.
593 Academic Press LTD, London UK. pp1-551.

594 Gibert, J., D.C. Culver, M.-J. Dole-Olivier, F. Malard, M.C. Christman, and L.
595 Deharveng. 2009. Assessing and conserving groundwater biodiversity: synthesis
596 and perspectives. *Freshwater Biology* 54:930-941.

597 Gibert, J., and L. Deharveng. 2002. Subterranean ecosystems: A truncated
598 functional biodiversity. *Bioscience* 52:473-481.

599 Gledhill, T., D.W. Sutcliffe, and W.D. Williams.1993. *British Freshwater Crustacea*
600 *Malacostraca: A key with ecological notes*. Freshwater Biological Association

601 Gooch J.L., and S.W. Hetrick. 1979. The Relation of Genetic Structure to
602 Environmental Structure: *Gammarus minus* in a KarstArea. *Evolution* 33:192-206.

603 Goricki, S., and P. Trontelj. 2006. Structure and evolution of the mitochondrial control
604 region and flanking sequences in the European cave salamander *Proteus*
605 *anguinus*. *Gene* 378:31-41.

606 Hahn, H.J. 2006. The GW-Fauna-Index: A first approach to a quantitative ecological
607 assessment of groundwater habitats. *Limnologica* 36:119-137.

608 Hahn, H.J., and A. Fuchs, 2009. Distribution patterns of groundwater communities
609 across aquifer types in south-western Germany. *Freshwater Biology* 54:848-860.

610 Hancock, P.J., and A.J. Boulton. 2009. Sampling groundwater fauna: efficiency of
611 rapid assessment methods tested in bores in eastern Australia. *Freshwater*
612 *Biology* 54:902-917.

613 Holsinger J.R.1988. Troglobites: The evolution of cave dwellin organisms. *American*
614 *Scientist* 76:146-153.

615 Humphreys W.F. 2009. Hydrogeology and groundwater ecology: Does each inform
616 the other? *Hydrogeology Journal* 17:5-21

617 Jaume, D., and W.F. Humphreys. 2001. A new genus of epacteriscid calanoid
618 copepod from an anchialine sinkhole in northwestern Australia. *Journal of*
619 *Crustacean Biology* 21:157–169.

620 Knight, L.R.F.D. and T. Gledhill. 2010. The discovery of *Microniphargus leruthi*
621 Schellenberg, 1934 (Crustacea: Amphipoda: Niphargidae) in Britain and its
622 distribution in the British Isles. *Zootaxa* 2655:52-56.

623 Kristjansson B.K., and J. Svavarsson. 2007. Sub-glacial refugia in Iceland enabled
624 groundwater amphipods to survive glaciations. *American Naturalist* 170:292-296.

625 Larned S.T. 2012. Phreatic groundwater ecosystems: research frontiers for
626 freshwater ecology. *Freshwater Biology* 57:885-906.

627 Lefébure, T., C.J. Douady, M. Gouy, P. Trontelj, J. Briolay, and J. Gibert. 2006.
628 Phylogeography of a subterranean amphipod reveals cryptic diversity and
629 dynamic evolution in extreme environments. *Molecular Ecology* 15:1797–1806.

630 Lefébure, T., C.J. Douady, F. Malard, and J. Gibert. 2007. Testing dispersal and
631 cryptic diversity in a widely distributed groundwater amphipod (*Niphargus*
632 *rhenorhodanensis*) *Molecular Phylogenetics and Evolution* 42:676–686.

633 Lesnoff, M., Lancelot, R. (2012). aod: Analysis of Overdispersed Data. R package
634 version 1.3, URL <http://cran.r-project.org/package=aod>

635 Malard, F., and F. Hervant. 1999. Oxygen supply and the adaptations of animals in
636 groundwater. *Freshwater Biology* 41:1-30.

637 Malard, F., C. Boutin, A.I. Camacho, D. Ferreira, G. Michel, B. Sket, and F. Stoch.
638 2009. Diversity patterns of stygobiotic crustaceans across multiple scales in
639 Europe. *Freshwater Biology* 54: 756-776.

640 Martin, P., C. De Broyer, F. Fiers, G. Michel, R. Sablon, and K. Wouters. 2009.
641 Biodiversity of Belgian groundwater fauna in relation to environmental conditions.
642 *Freshwater Biology* 54:814-829.

643 Mauclaire, L. and J. Gibert. 2001. Environmental determinants of bacterial activity
644 and faunal assemblages in alluvial riverbank aquifers. *Archiv für Hydrobiologie*
645 152:469-487.

646 Maurice, L. and J. Bloomfield. 2012. Stygobitic invertebrates in groundwater — A
647 review from a hydrogeological perspective. *Freshwater Reviews* 5:51-71.

648 Maurice, L.D., T.C. Atkinson, J.A. Barker, J.P. Bloomfield, A.R. Farrant, and A.T.
649 Williams. 2006. Karstic behaviour of groundwater in the English Chalk. *Journal of*
650 *Hydrology*. 330: 63-70.

651 McInerney C.E., L. Maurice, A.L. Robertson, L.R.F.D. Knight, J. Arnsheidt, C.
652 Venditti, J.S.G. Dooley, T. Mathers, S. Matthijs, K. Eriksson, G. Proudlove and B.
653 Hänfling (2014) The ancient Britons: Groundwater fauna survived extreme
654 climate changes over tens of millions of years across NW Europe. *Molecular*
655 *Ecology* 23: 1153-1166.

656 Moldovan O.T., I.N. Meleg, and A. Persoiu. 2012. Habitat fragmentation and its
657 effect on groundwater populations. *Ecohydrology* 5:445-452.

658 Notenboom, J., 1991. Marine regressions and the evolution of groundwater dwelling
659 amphipods (crustacea). *Journal of Biogeography* 18:437-454.

660 Paran F., F. Malard J. Mathieu, M. Lafont, D.M. P. Galassi, and P. Marmonier. 2005.
661 Distribution of groundwater invertebrates along an environmental gradient in a
662 shallow water-table aquifer. Pages 99-105 in J. Gibert (editor) *World*
663 *Subterranean Biodiversity, Proceedings of an International Symposium* (Ed. J.
664 Gibert) University Claude Bernard, Lyon, France

665 Price M. 1996. *Introducing Groundwater*, 2nd edn. Chapman and Hall, London. pp
666 70-87.

667 Proudlove, G.S., P.J. Wood, P.T. Harding, D.J. Horne, T. Gledhill, and L.R.F.D.
668 Knight.2003. A review of the status and distribution of the subterranean aquatic
669 Crustacea of Britain and Ireland. *Cave and Karst Science* 30:51-74

670 R Core Team (2013). *R: A language and environment for statistical computing*.
671 R Foundation for Statistical Computing, Vienna, Austria. URL [http://www.R-proje](http://www.R-project.org/)
672 [ct.org/](http://www.R-project.org/).

673 Robertson, A. L., J.W.N. Smith, T. Johns, and G.S. Proudlove. 2009. The distribution
674 and diversity of stygobites in Great Britain: an analysis to inform groundwater
675 management. *Quarterly Journal of Engineering Geology and Hydrogeology*
676 42:359-368.

677 Rukke, N.A. 2002. Effects of low calcium concentration on two common freshwater
678 crustaceans, *Gammarus lacustris* and *Astacus astacus*. *Functional Ecology*
679 16:357–366.

680 Scheidleder, A., and P. Visser, 2012. Comparative Study of Pressures and
681 Measures in the Major River Basin Management Plans: Drinking water protected
682 areas: Task 3c – Summary note on drinking water safeguard zones. European
683 Commission. 3-4.

684 Schminke, H,K. 1974. Mesozoic Intercontinental Relationships as Evidenced by
685 Bathynellid Crustacea (Syncarida : Malacostraca). *Systematic Biology* 23:157-
686 164.

687 Sket, B. 2004. Biodiversity in hypogean waters page 709. *In: J. Gunn (editor)*
688 *Encyclopedia of Caves and Karst Science*. Fitzroy Dearborn, New York,

689 Smith, H., P.J. Wood, and J. Gunn 2003. The influence of habitat structure and flow
690 permanence on invertebrate communities in karst spring systems. *Hydrobiologia*
691 510: 53-66.

692 Stanford, J.A., and J.V. Ward. 1993. An ecosystem perspective of alluvial rivers;
693 connectivity and the hyporheic corridor. *Journal of the North American*
694 *Benthological Society* 12:48-60.

695 Stock, J., 1980. Regression model evolution as exemplified by the genus
696 *Pseudoniphargus* (Amphipoda). *Bijdragen tot de Dierkunde* 50:105–144.

697 Stoch, F., and D.M.P. Galassi, 2010. Stygobiotic crustacean species richness: a
698 question of numbers, a matter of scale. *Hydrobiologia* 653:217-234.

699 Strayer, D.L., 1994. Limits to biological distributions in groundwater. *in* J. Gibert, D.
700 Danielopol and J. Stanford. (editors) *Groundwater Ecology*. Academic Press
701 Limited, London. U.K. pp 287-310.

702 ter Braak, C.J.F and P. Smilauer. 1998. CANOCO reference manual and user's
703 guide to Canoco for windows: software for canonical community ordination
704 (version 4). Wageningen.

705 Trontelj, P., C.J. Douady, C. Fiser, J. Gibert, S. Goricki, T. Lefebure, B. Sket, and V.
706 Zaksek, V., 2009. A molecular test for cryptic diversity in ground water: how large
707 are the ranges of macro-stygobionts? *Freshwater Biology* 54:727-744.

708 Ward, J.V., and M.A. Palmer. 1994. Distribution patterns of interstitial freshwater
709 meiofauna over a range of spatial scales, with emphasis on alluvial river-aquifer
710 systems. *Hydrobiologia*. 287:147-156.

711

712

713

714 FIGURE LEGENDS

715 Figure 1: Distribution of groundwater fauna sampling stations and hydro-units within
716 the study region in the south-west of England.

717 Figure 2: Stygobitic species presence and community composition at sites within the
718 study region in the south-west of England. The size of the circles reflects the
719 number of species recorded.

720 Fig. 3: CCA ordination: Ordination by canonical correspondence analysis of
721 stygobiont species and stygophile data with significant environmental variables
722 (identified using the forward selection procedure) from boreholes, wells and springs
723 in SW England (Sept 2009-Nov 2011).

724

725 Table 1: Classification and description of the 5 hydro-units in the study area

Hydro-unit (no.of sites)	Geology	Main geologies	% Area coverage	Flow	Permeability (relative)
1 (10)	Granular	Quaternary & Palaeogene unconsolidated material	6.3	intergranular	Low-high
2 (37)	Igneous & metamorphic	Granite, basalt, spillite & gneiss & mica schists	8.6	Fracture	Low-mid
3 (57)	Mudstones & siltstones	Carboniferous, Triassic & Jurassic mudstones	51.6	Fracture and intergranular	Very low
4 (55)	Sandstones	Permo-Triassic, Jurassic & Devonian sandstones	17.2	Fracture and intergranular	Mid-high
5 (63)	Carbonate rocks	Chalk, Carboniferous & Devonian limestones	16.4	Karstic (solutional fissures and conduits)	Low-high

726

727

728 Table 2: Means and standard deviations of water chemistry parameters for each hydro-
 729 unit. Kruskal-wallis results indicate where determinands were significantly different
 730 between hydro-units.

731

Hydro-Unit	Statistic	EC μScm ⁻¹	DO mg/l	Temp °C	pH	DOC mg/l	NH ₄ mg/l	Cl mg/l	NO ₃ mg/l	SO ₄ mg/l	Ca mg/l	Mg mg/l	K mg/l	Na mg/l
1	n	8	8	8	8	8	8	8	8	8	8	8	8	8
	Mean	370.25	3.65	13.71	7.04	3.63	0.52	9.59	7.46	10.29	47.51	3.53	3.76	16.00
	Std Dev.	199.84	3.58	0.55	0.73	1.19	0.74	3.90	4.67	10.43	26.88	3.12	3.51	5.57
2	n	37	14	23	36	32	13	37	30	32	32	32	27	33
	Mean	276.14	8.91	9.65	6.35	1.41	0.02	8.86	4.66	4.42	25.07	12.33	4.95	20.46
	Std Dev.	193.32	3.41	1.36	0.76	1.72	0.01	7.18	4.85	3.36	12.37	13.98	6.11	14.47
3	n	47	29	29	45	47	29	46	41	47	46	46	45	45
	Mean	349.70	5.53	10.38	6.56	3.65	0.07	11.33	4.10	8.69	30.13	13.68	5.14	23.43
	Std Dev.	198.71	4.32	1.54	0.51	9.38	0.18	6.01	5.99	8.52	16.80	9.93	6.49	13.73
4	n	37	24	26	37	39	6	40	26	37	38	38	34	38
	Mean	423.89	5.90	10.60	6.81	11.52	0.82	12.14	9.07	11.04	37.68	7.40	5.92	18.22
	Std Dev.	245.74	3.89	1.44	0.75	15.48	1.63	6.77	8.30	12.87	27.10	5.21	7.63	9.23
5	n	54	25	39	54	54	17	55	53	52	53	52	43	53
	Mean	582.96	5.69	11.50	7.09	12.37	0.10	11.81	9.35	6.76	55.64	4.51	4.08	15.79
	Std Dev.	133.64	3.14	0.82	0.43	31.16	0.11	6.02	5.52	4.49	21.89	4.25	3.21	8.82
Kruskal Wallis Test	H Df p	48.54 4 <0.001	11.36 4 <0.001	22.37 4 <0.001	30.49 4 <0.001	53.51 4 <0.001	N/A	14.25 4 <0.01	38.66 4 <0.001	18.56 4 <0.01	44.73 4 <0.001	36.24 4 <0.001	5.51 4 0.239	13.44 4 <0.01

732 N/A – insufficient data (majority of values below detection limit).

733

734

735 Table 3: Abundance and composition of groundwater taxa (stygo-bionts and
 736 stygo-philic) in boreholes (BH), wells and springs from the five hydro-units.
 737

Attributes of sample sites	Total	Sites sampled (no.)		
		BHs	Wells	Springs
No. of sites sampled:	221	124	56	41
<u>No. of sites in each hydro-unit (Hu)</u>				
Granular (Hu1)	10	9	1	0
Igneous/metamorphic (Hu2)	37	12	12	13
Mudstone/siltstone (Hu3)	56	28	19	9
Sandstone (Hu4)	55	34	11	10
Carbonate (Hu5)	63	41	13	9
No. of sites with GW fauna	155	70	51	34
Sites with no GW fauna* ¹	66	54	5	7
No. of sites with stygo-bionts	75	34	23	18

Taxa groups & species	Total Abundance	Percentage occurrence from all sites				Hydro-units present
Stygo-bionts	465	33.8	27.4	40.4	43.9	All
<u>Crustacea, Malacostraca, Eumalacostraca</u>						
<u>Peracarida</u>						
<u>Amphipoda</u>						
<u>Niphargidae</u>						
<i>Niphargus kochianus</i>	181	10.8	12.9	12.3	2.5	3, 5
<i>Niphargus fontanus</i>	28	5.4	4.8	7.0	4.9	1, 5
<i>Niphargus aquilex</i>	96	10.4	8.1	12.3	14.6	1, 2, 3, 4, 5
<i>Niphargus glenniei</i>	79	8.6	2.4	12.3	22.0	2, 3, 4, 5
<i>Microniphargus leruthi</i>	10	2.7	2.4	3.5	2.5	3, 4, 5
<u>Crangonyctidae</u>						
<i>Crangonyx subterraneus</i>	60	5.9	6.5	3.5	7.3	4, 5
<u>Isopoda</u>						
<u>Asellidae</u>						
<i>Proasellus cavaticus</i>	10	2.3	4.0	0	0	5
<u>Syncarida</u>						
<u>Bathynellacea</u>						
<u>Bathynellidae</u>						

<i>Antrobathynella stammeri</i>	1	0.5	0.8	0	0	4
Stygophiles	5318	63.5	44.4	86.0	90.2	All
<u>Crustacea, Maxillopoda</u>						
Copepoda	4181	47.3	26.6	70.2	78.0	1, 2, 3, 4, 5
<u>Crustacea, Ostracoda</u>	745	23.4	8.1	38.6	48.8	1, 2, 3, 4, 5
<u>Arthropoda, Arachnida, Acari, Trombidiformes</u>						
Hydrachnidae	182	16.2	10.5	21.1	26.8	1, 2, 3, 4, 5
<u>Annelida, Clitellata</u>						
Oligochaeta* ²	210	10.4	12.9	12.3	-	2, 3, 4, 5

738 *¹ Other freshwater (epigean) or terrestrial taxa may have been recorded at these sites

739 *² Not included where recovered from spring sources, due to presence of epigean species

740

741 Table 4: Summary of odds ratios from generalized linear models selected to describe
742 relationships between presence/absence data, hydro-units (Hu) and water chemistry.
743 (The level of significance reported in each model is represented by: * 0.05, ** 0.01,
744 ***0.001).

Odds ratios calculated relative to:

Hydro-unit 5 (carbonate)

Sample type 1
(Boreholes)

Generalized linear model (simplest model selected based on comparison of all model permutations)	Determinand (value)	Granular material (Hu1)	Igneous / metamorphic (Hu2)	Mudstone / siltstone (Hu3)	Sandstone (Hu4)	Sample type 2 (wells)	Sample type 3 (springs)
Pres/Abs~Hydro-unit	all data	0.84	0.80	0.12***	0.21***		
$\chi^2 = 33.8, df=5, p<0.001$							
	dissolved oxygen	2.00	1.00	0.16**	0.26*		
	nitrate	N/A	0.67	0.12***	0.28*		
	temperature	2.38	1.48	0.15**	0.23*		
Pres/Abs~Hydro-unit + value	calcium	0.97	0.98	0.18**	0.27**		
	chloride	0.93	0.63	0.11**	0.20**		
	pH	0.96	0.56	0.08***	0.16***		
Pres/Abs~Hydro-unit + sample type	Dissolved organic carbon	1.12	0.65	0.10***	0.21***	2.34*	3.01*
	elec. conductivity	1.09	0.59	0.10***	0.22**	2.19	3.07*
	magnesium	0.94	0.43	0.10**	0.19***	1.83	3.22*
	potassium	0.91	0.30*	0.07***	0.14***	1.77	4.52*
	sodium	1.00	0.49	0.12***	0.21**	1.92	3.20*
	sulphate	1.89	0.49	0.09***	0.19**	1.86	3.51*

746 N/A – excluded from analysis because insufficient reliable nitrate data available

747 Level of significance in model * 0.05, ** 0.01, ***0.001

748

749

750

751

752

753

754

755

756 Table 5: Partialled CCA results showing percentage of variation in species data explained
757 by environmental variables; hydro-units & groundwater chemistry, together and in isolation.

758

CCA	Constraint on Species data	Variable partialled out (covariable)	Sum of all canonical eigen-values (Ev)	Percentage variation (Ev x 100/4.998)	Fraction of variation explained by
CCA1	Chemistry	none	0.429	8.58%	-
CCA2	Hydro-units	none	0.628	12.57%	-
CCA3	Hydro-units	Chemistry	0.251	5.02%	Chemistry
CCA4	Chemistry	Hydro-units	0.376	7.52%	Hydro-units
Sum of all eigenvalues in a species only CA			4.998		
Overall amount of variation due to environmental variables				16.11%	CCA1 + CC4
Variation due to hydro-units & water chemistry together				3.56%	CCA1 –CCA3
Variation unexplained by hydro-units &/or water chemistry				83.89%	100 – (CCA1 + CCA4)

759