

## Loughborough University Institutional Repository

---

# *External heating garments used post warm-up improve upper-body power and elite sprint swimming performance*

This item was submitted to Loughborough University's Institutional Repository by the/an author.

**Citation:** WILKINS, E.L. and HAVENITH, G., 2017. External heating garments used post warm-up improve upper-body power and elite sprint swimming performance. *Proceedings of the Institution of Mechanical Engineers Part P: Journal of Sports Engineering and Technology*, 231(2), pp.91-101.

**Additional Information:**

- This paper was accepted for publication in *Proceedings of the Institution of Mechanical Engineers, Part P: Journal of Sports Engineering and Technology*. The definitive version is available at: <http://dx.doi.org/10.1177/1754337116650322>

**Metadata Record:** <https://dspace.lboro.ac.uk/2134/21221>

**Version:** Accepted for publication

**Publisher:** SAGE Publications / © IMechE

**Rights:** This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Please cite the published version.

1 External Heating Garments used Post Warm-Up Improve Upper-Body Power and Elite  
2 Sprint Swimming Performance

3 **Emma L Wilkins<sup>1,2</sup> and George Havenith<sup>1</sup>**

4

5 <sup>1</sup>Environmental Ergonomics Research Centre, Loughborough Design School, Loughborough  
6 University, Leicestershire, UNITED KINGDOM;

7 <sup>2</sup>School of Sport, Exercise and Health Sciences, Loughborough University, Leicestershire, UNITED  
8 KINGDOM;

9

10 **Running head: Muscle heating garment use in sprint swimming**

11

12

13

14

15

16 Address for correspondence:

17 George Havenith,

18 Environmental Ergonomics Research Centre,

19 James France Building,

20 Loughborough University,

21 Loughborough, Leicestershire, LE11 3TW,

22 +44 1509 223031,

23 G.Havenith@lboro.ac.uk

24

## 25 ABSTRACT

26

27 The aim of this study was to determine the effects of using an electrical heating garment during  
28 a 30-minute recovery period after a standardized swimming warm-up on subsequent swimming  
29 performance and upper-body power output. On two occasions, eight male and four female elite  
30 competitive swimmers completed a standardized swimming warm-up, followed by a 30-minute  
31 passive recovery period before completing maximal plyometric press-ups and a 50m Freestyle  
32 swim. Plyometric press-ups determined starting strength (SS), peak force (PF) and peak  
33 concentric power (PCP). During the recovery period, participants wore tracksuit bottoms and (i)  
34 a standard tracksuit top (CON) or (ii) jacket with integrated electric heating elements (HEAT).  
35 The overall results demonstrated a trend of a relevant (>0.4%) improvement in the 50m  
36 Freestyle performance of 0.83% ( $P = 0.06$ ) in HEAT vs. CON. In male participants,  
37 performance in the 50m Freestyle significantly improved by 1.01% (CON  $25.18 \pm 0.5s$  vs.  
38 HEAT  $24.93 \pm 0.4s$ ;  $P < 0.05$ ), whereas female participants only showed a trend for an  
39 improvement of 0.38% ( $29.18 \pm 0.5s$  vs.  $29.03 \pm 1.0s$ ;  $P = 0.09$ ), in HEAT compared with  
40 CON, though statistical power for the latter test was low. Male participants' starting strength,  
41 peak force and peak concentric power were  $16.5 \pm 13\%$ ,  $18.1 \pm 21\%$  and  $16.2 \pm 21\%$  greater,  
42 respectively, in HEAT compared with CON (all  $P < 0.01$ ). In conclusion, external heating of the  
43 upper body between completion of the warm-up and performance through the utilization of an  
44 electrically heated jacket improves plyometric press-up power output and force production, as  
45 well as sprint swimming performance in males. This provides justification for future  
46 enhancement opportunities in sporting performance through the utilization of external heating  
47 systems. Optimization of the heating system for specific sports is required.

48

49

50 **Key Words:** MUSCLE TEMPERATURE, SWIMMERS, CLOTHING, PASSIVE HEATING,  
51 GENDER

## 52 1. INTRODUCTION

53  
54 The importance of warming-up on the enhancement of exercise performance is well established (1). Its  
55 impact on subsequent performance is dependent on the intensity and duration of a competition event  
56 and on the recovery duration between the warm-up and the competitive event (2). One of the major  
57 contributing factors to a heightened performance is an increase in muscle temperature ( $T_m$ ), with  
58 increases of 3°C - 4°C shown following an active warm-up (3). Not only can  $T_m$  maintenance be pivotal  
59 between the warm-up and event, but also between multiple races at an event (4). Due to time  
60 constraints, a swimmer may not be able to change into a dry racing suit between races; meanwhile,  
61 remaining in the wet suit increases body heat loss and speeds  $T_m$  cooling.

62 A warm-up induced rise in  $T_m$  results in a number of beneficial physiological effects (5, 6, 7, 8), ranging  
63 from increased anaerobic metabolic capacity, increased nerve conduction rates in both the central and  
64 peripheral nerves, and increased speed of muscle contractions, to adjustments in muscle sensitivity and  
65 calcium production. All these together lead to significantly improved muscle function, force and power  
66 production; and subsequently, to improved performance (5, 6), whether  $T_m$  is raised as a result of  
67 exercise or passive heating (9).

68  
69 Most of the existing literature advocates the benefits of increased  $T_m$  on short duration events (<5  
70 minutes), which have a greater dependence on high levels of power production (5, 10, 11). Bergh and  
71 Ekblom (10) revealed a 5% increase in power output, jumping, and sprinting performances for each 1°C  
72 increase in  $T_m$ , between muscle temperatures of 30 and 39°C, via cooling and warming experiments.  
73 Faulkner *et al.* (6) observed a 9% increase in peak power output per-degree-centigrade elevation in  $T_m$ .  
74 Faulkner *et al.* (6, 12), studying cycling sprint performance, reported on the problems with dropping  $T_m$   
75 occurring when the warm-up and race are separated by a period in which the athlete is inactive. For

76 swimmers, such an inactive period between the warm-up and race is common. In national and  
77 international competitions, swimmers must report to the call room 20-minutes prior to their race, with  
78 most swimmers completing their warm-up up to 45-minutes prior to racing (2). The impact of such a  
79 delay was studied by Zochowski *et al.* (2) who observed a ~1.4% better 200m freestyle performance of  
80 national standard swimmers after a 10-minute post warm-up recovery/delay, in comparison to a 45-  
81 minute recovery/delay period. Similarly, West *et al.* (7) observed 200m swim times to be  $1.86 \pm 1.37s$   
82 better when swam within 20-minutes of the warm-up in comparison to 45 minutes, resulting in a  
83 difference of a 1.5% improvement in performance. The predicted higher  $T_m$  in the shorter recovery  
84 periods (6, 12) is assumed to be the underlying cause for these observed performance enhancements.

85

86 Such an observed improvement of 1.5% in performance is of great significance to an elite swimmer.  
87 According to Pyne *et al.* (13), swimmers can substantially increase their chances of medaling by  
88 improving performance by as little as 0.4%, demonstrated at the 2012 Olympics where the bronze  
89 medal position and 4<sup>th</sup> position were separated by just 0.09% (0.02 seconds) and 0.25% (0.07 seconds)  
90 respectively in the men's and women's 50m Freestyle.

91

92 With studies finding a significant deterioration in high-power performances of short duration (< 5  
93 minutes) after prolonged periods between the warm-up and competing, development of methods to keep  
94 a raised  $T_m$  during this recovery period are crucial (3, 6). The main focus in the development of such  
95 methods has been on heated trousers. Faulkner *et al.* (6, 12) demonstrated the benefits of external  
96 heating (heated trousers) between warm-up completion and racing in sprint cycling, achieving a  $\sim 1^\circ C$   
97 higher  $T_m$  and a concomitant 9% increase in peak power (6, 12) with a 4% increase in mean power (12),  
98 compared to wearing a normal track suit in the 30-minute recovery period.

99

100 Since 90% of maximal freestyle velocity is produced by the arms, with only 10% propulsion from the  
101 legs (14), upper-body heating for swimmers is more relevant, thus the focus for this sport should be on a  
102 heating jacket rather than trousers. Studying national junior swimmers, McGowan *et al.* (15) found that  
103 adding the wearing of heated jackets to dryland-based exercise circuits between warm-up completion  
104 and racing (30 minutes) further improved the 100m Freestyle swimming performance above the dryland  
105 exercises alone, though the heated jackets on their own did not increase performance (15). Given that  
106 performances and the impact of performance-altering interventions often relate to the level of athletes  
107 investigated and the distance covered, it is unclear whether McGowan *et al.*'s results would also  
108 translate into elite senior swimmers and shorter sprint events.

109

110 Based on these considerations, it was felt that an additional study investigating the impact of heated  
111 garments, and more specifically heated jackets, in the recovery period of elite senior swimmers (rather  
112 than McGowan's juniors) was relevant for the evaluation of such techniques. Furthermore, where  
113 McGowan *et al.* (15) used the 100m freestyle, it was considered that a 50m freestyle sprint would be  
114 most relevant to test the impact of muscle heating, given that the biggest impact of a heating procedure  
115 is expected in a short burst of high-power exercise. An improvement would be considered relevant if  
116 higher than 0.4%, based on the work of Pyne *et al.* (13). Apart from directly investigating sprint  
117 performance, upper-body performance was also investigated as an additional performance measure  
118 (16), to see whether the hypothesized higher  $T_m$  due to the application of the heating jackets would  
119 produce a measurable effect of instantaneous upper-body power in short duration. Bench press exercise  
120 has been linked to arm-force production and better swimming times in water (16, 17), but utilizing this  
121 exercise was not technically feasible in this setup. Plyometric press-up power output has also been  
122 linked to enhanced swimming training and performance (18), be it less directly. Given the general link  
123 between plyometric press-up and upper-body power, and the link between the latter and swimming  
124 performance, this method was arbitrarily chosen as a secondary measure that could form the basis for  
125 any observed improvement in sprint swimming performance.

## 126 2. METHOD

127 **Experimental Approach to the Problem**

128

## 129 2.1 PARTICIPANTS

130 Twelve participants, eight elite male swimmers (aged =  $21 \pm 1.8$  yr, height =  $1.88 \pm 0.06$ m, body mass  
131 =  $87.6 \pm 7.65$  kg, FINA points (2014) =  $684 \pm 56$ ; mean  $\pm$  SD) and four elite female swimmers (aged =  
132  $20 \pm 1.7$  yr, height =  $1.72 \pm 0.09$ m, body mass =  $66.9 \pm 10.14$  kg, FINA points =  $651 \pm 10$ ; not  
133 controlled for menstrual cycle) volunteered to participate in this study. An elite swimmer is defined as  
134 an athlete that is of adult age who is close to or has already reached their top performances, competing  
135 regularly at the key national- or international-level competitions (19). Sample size was defined using  
136 the model of Hopkins (20) (change in mean in a crossover study), based on the standard deviation of  
137 non-tapered performance times and the smallest worthwhile enhancement in performance of 0.4% (13).  
138 This analysis indicated the need for 8 participants. Four female participants were added to the group of  
139 eight male participants to investigate possible gender impacts on the results. Due to logistical reasons,  
140 unfortunately, a complete sample of 8 females, needed for appropriate power to analyze gender data  
141 separately, was not achieved. Nevertheless, the data are included here and presented with consideration  
142 of the low statistical power. The 50m Freestyle personal best times for male and female participants of  
143 this study were  $23.83 \pm 0.76$  seconds and  $27.15 \pm 0.66$  seconds, respectively (mean  $\pm$  SD). All  
144 participants performed at least seven swimming sessions per week ( $16.7 \pm 1.6$  h  $\text{wk}^{-1}$ ) along with 2-3  
145 land-based sessions ( $5.9 \pm 0.7$  h  $\text{wk}^{-1}$ ), and had  $13.3 \pm 2.7$  years of practice which indicates expert skill  
146 (21). Participants were informed of the benefits and risks of the study prior to giving their written  
147 informed consent to participate in the study. Participants completed a general health-screen  
148 questionnaire and were all non-smokers and free from injury. The study was carried out during the  
149 swimmers' competitive season to ensure a high state of physical training. The study was approved by  
150 the Loughborough University Ethical Advisory Committee.

## 151 2.2 STUDY OVERVIEW



152 Prior to the experimental trials, participants were familiarized with the testing protocol, as well as  
153 measurements and exercise testing. Also preceding the experimental trials, participants completed a  
154 two-week pilot study assessing plyometric press-ups as a performance measure, in order to minimize  
155 the learning effect during the course of the study. Within-subject coefficient of variation (CV) %  
156 calculations indicated starting strength (CV%=8.26) and peak force (CV%=3.44) to have moderate to  
157 very high test-retest reliability, in agreement with Hogarth *et al.* (220), whereas peak concentric power  
158 (W) (CV%=10.78) demonstrated a CV just above the analytical goal of  $\geq 10\%$  (23).

159

160

161 Participants visited the swimming pool for two testing sessions. Each time, they completed a 30-minute  
162 standardized swimming warm-up, followed by a period of 30-minute passive seated recovery,  
163 simulating the time between finishing the warm-up and racing. During the 30-minute seated recovery,  
164 participants underwent one of two conditions: wearing either the standardized jacket (CON) or the  
165 heated jacket (HEAT) (detailed below) followed by, after removal of the clothing, four plyometric  
166 press-ups and a maximum long-course 50m Freestyle. A repeated-measures study design was utilized,  
167 with each swimmer completing both a control and intervention trial, separated by seven days. Trial  
168 conditions were performed in a balanced order and took place at the same time of day (~14:00), aiming  
169 to minimize circadian variations effects on performance. Participants completed their performance  
170 measures individually to avoid any external influences. Twenty-four hours prior to testing, participants  
171 were asked to refrain from caffeine and alcohol consumption, as well as any strenuous exercise. Passive  
172 recovery was carried out in a temperature-controlled room ( $20.0 \pm 0.2^{\circ}\text{C}$ ), to simulate competition cool  
173 rooms and ensure consistent conditions across tests. Warm-up and swimming tests were carried out in  
174 an Olympic standard 50m swimming pool (Pool water temperature  $27.6 \pm 0.1^{\circ}\text{C}$ , Air temperature  $23.4$   
175  $\pm 0.1^{\circ}\text{C}$ , Humidity  $55.8 \pm 1.4\%$ ) at Loughborough University.

176

177 **Procedures**

178 Participants arrived at the pool after a typical competition-day meal at least two hours prior to testing  
179 (repeated over trials). Upon arrival, participants had their height (Esca, Birmingham, United Kingdom)  
180 and body mass (M) (Esca 770, Vogel & Halke, Hamberg, Germany) recorded, from which body surface  
181 area ( $A_D$ ) was estimated and surface-to-mass ratios ( $A_D/M$ ) of the subjects were calculated. Body fat  
182 percentage data were based on seven-point skinfold measurements. Participants entered the  
183 temperature-controlled room and remained seated for a 15-minute stabilization period. All participants  
184 wore a standardized tracksuit: a single layer of uninsulated nylon material consisting of trouser bottoms  
185 and a zip-up top. During this time, they were familiarized with the trial procedure. Following the  
186 stabilization period, a baseline skin thermal image (FLIR i7, Flir Systems, Wilsonville, USA) of  
187 participants in their swimsuit was captured from a distance of 3m, in anatomical position with palms  
188 facing forward, along with measurements of tympanic temperature (TT) (Braun ThermoScan PRO  
189 4000, Welch Allyn, Kaz, USA), heart rate (HR) (Polar FT1, Polar Electro Oy, Kempele, Finland),  
190 thermal comfort (TC) and thermal sensation (TS) (24).

191

192 Participants then completed a standardized heart rate (HR)-monitored swimming warm-up, with the HR  
193 noted after completion of the 4 x 50m sprinted bursts. The warm-up is a standardized warm-up as  
194 described by West et al. (7), with the 4 x 50m altered to make it more sprint-focused. The warm-up  
195 entailed: 400 m Freestyle, 200 m Pull, 200 m Kick, 200 m Drill (Fins), 200 m Individual Medley, 4 ×  
196 50m Freestyle: (1) Push 15m u/w fly kick, (2) 15m spin drill, (3) dive 15m race pace, (4) dive 25m race  
197 pace (HR measured), 200 m easy.

198 Promptly after the completion of the warm-up, skin thermal imaging, heart rate, thermal comfort and  
199 thermal sensation were recorded as described above. Participants then remained seated for 30 minutes  
200 in the temperature-controlled room, simulating a call-room marshalling period. Participants wore a  
201 standard pair of tracksuit bottoms, long-sleeve top and one of two types of jackets that made up the  
202 intervention: 1) control (CON) where participants wore a standardized tracksuit jacket, or 2) heated  
203 jacket (HEAT) where participants wore a jacket with integrated heated elements (Powerlet rapidFIRE

204 Proform Heated Jacket Liner, Warren, USA). When unheated, both jackets had similar insulation as  
205 measured on a thermal manikin (25). The heated jacket was selected based on market research to find  
206 the best coverage of the torso and arms with heating elements. The heated elements targeted the major  
207 muscle groups: pectoralis major, latissimus dorsi, tricep brachii, and covered the lower deltoids (26)  
208 (Fig. 1). The heating elements were powered by 12v 10amp power transformers powering the jacket to  
209 full capacity at 105watts, with the elements reaching temperatures of  $\sim 50^{\circ}\text{C}$  (lower on the skin contact).  
210 The jacket's stretch panels allow maximum heat transfer, as the material stays in close contact to the  
211 body, reducing convection whilst permitting movement. Over the duration of the 30-minute period,  
212 measurements of thermal comfort and thermal sensation were recorded every 5 minutes.

213 Subsequently, tracksuit garments were removed and a further skin thermal image was captured in their  
214 swimsuit alone. Thermal images were analyzed using ThermoCAM Researcher Software (Flir,  
215 Wilsonville, USA) to measure mean skin temperature ( $T_{sk}$ ) of the upper body (torso and upper arms)  
216 using the freeform tool. Muscle temperatures ( $T_m$ ) were estimated from mean skin temperature as:  $T_m$   
217  $= 1.02T_{sk} + 0.89$  ( $r^2=0.98$ ), based on work by De Ruiter *et al.* (27). Tympanic temperature, heart rate,  
218 thermal comfort and thermal sensation were also recorded.

219 Each participant then performed four separate maximal-effort plyometric press-ups (without the heating  
220 garment), with  $\sim 10$ s rest in between, on a force platform (400S Force Plate, Fittech, Skye, Australia,  
221 sampled at 600Hz) whereby kinetic data were collected and analyzed using Ballistic Measurement  
222 System Software. After the force platform was reset, participants were instructed to place their hands at  
223 a self-selected width, with elbows straight. Male participants performed a regular press-up with feet  
224 together, whereas female participants performed bent-knee press-ups, a lower-intensity press-up  
225 variation (28). Succeeding a three-second count down, participants performed the countermovement  
226 action of the plyometric press-up as quickly as possible, aiming for maximal height of trunk elevation.  
227 Force data were analyzed to determine Starting Strength (SS, also called 'maximal rate of force  
228 development', calculated as the steepest slope of the force time curve), Peak Force (PF, highest  
229 measured value) and Peak Concentric Power (PCP). After completion, participants placed the respective

230 jacket back on and prepared themselves to perform a 50m Freestyle time trial at maximum effort (~2  
231 minutes after completing press-ups).

232

233 For the swim trial, a starter system (HS-200 Horn Start, Daktronics, Inc., Brookings, SD, USA) was  
234 used to replicate the signal used in competitions. Participants began their swim (without a race suit)  
235 from the blocks (Omega® OSB11, Swiss Timing, Switzerland) to simulate race conditions, and the 25m  
236 split, stroke rates (at ~20m & 40m, i.e. Stroke Rate 1 and Stroke Rate 2) and total stroke count over the  
237 whole distance were recorded. An official electronic timing system with an accuracy of 1/1000s  
238 (Omega Ares 21, Swiss Timing, Switzerland) was used to determine the overall swim time, with 25m  
239 splits taken by the coach using a stopwatch (Fastime 9, Pyramid Technologies, Meriden, CT).  
240 Immediately after completion, the HR was measured, followed by thermal comfort, thermal sensation  
241 and rating of perceived exertion (RPE) using Borg's 15-point-scale (29).

242

### 243 **Statistical Analyses**

244 All statistical tests were processed using IBM SPSS Statistics Software Version 22. The Shapiro-Wilk  
245 test of normality revealed the data were normally distributed. Participant characteristics were analyzed  
246 using an independent-samples T-test. Performance data were analyzed using a one-tailed, paired T-test  
247 based on the directional hypothesis.  $T_{sk}$ , TT and HR were analyzed (two tailed) using a one-way  
248 repeated measures ANOVA (Condition \* Time). RPE, TC and TS data among participants were  
249 analyzed using the Freidman Test. Significant effects were followed up with the Wilcoxon Signed-Rank  
250 Test, and the Kruskal-Wallis test was followed up by the Mann-Whitney *U* Test for between genders.  
251 The accepted level of significance was  $P < 0.05$ , with a trend level of  $0.05 < P < 0.1$  also being  
252 acknowledged. Data are presented as mean  $\pm$  SD. The 50m Freestyle performance was further analyzed  
253 using Hopkins' (20) published spreadsheet that used log transformation to estimate the effect of passive  
254 heating as the difference in the mean percent change between the experimental and control groups. The  
255 spreadsheet provided the precision of the estimate and the chances that the true effect was practically

256 beneficial or harmful at a 90% confidence limit. For calculations of the chances of benefit and harm, the  
257 value of 0.4% for the smallest worthwhile effect was used (13). Quantitative chances of benefit or harm  
258 were assessed qualitatively as follows: <1%, almost certainly not; 1-5%, very unlikely; 5-25%, unlikely;  
259 25-75%, possible; 75-95%, likely; 95-99, very likely; and >99%, almost certain.

### 260 3. RESULTS

#### 261 3.1 PARTICIPANT CHARACTERISTICS

262 The male participants were significantly taller than the female participants ( $1.88 \pm 0.06$  vs  $1.72 \pm 9$  m,  $P$   
263  $< 0.05$ ) and had a greater body mass ( $87.6 \pm 7.6$  vs  $66.9 \pm 10.1$  kg,  $P < 0.05$ ); thus, they had a  
264 significantly larger body-surface-area ( $2.1 \pm 0.1$  vs  $1.8 \pm 0.2$  m<sup>2</sup>,  $P < 0.05$ ) but lower body-surface-area  
265 to mass-ratio than the female participants ( $245 \pm 7 \cdot 10^{-4}$  vs  $269 \pm 14 \cdot 10^{-4}$  m<sup>2</sup> kg<sup>-1</sup>,  $P < 0.05$ ). The male  
266 participants had a significantly lower body fat percentage than the female participants ( $6.1 \pm 2.2$  vs  $21.0$   
267  $\pm 4.6\%$   $P < 0.05$ ).

268

#### 269 3.2 SWIMMING PERFORMANCE

270 When observing both male and female participants, a trend was shown in the 50m Freestyle time where  
271 HEAT performance was faster compared to that in CON by 0.83% ( $P = 0.06$ ), with a significant 1.06%  
272 improvement in the 25m split time ( $P < 0.05$ ) (Table 1). Eight of the twelve participants (six of the eight  
273 males and two of the four females) showed a clear improvement in swimming performance, improving  
274 by more than 0.4%- the smallest worthwhile enhancement in swimming (Fig. 2) (13). Stroke rate 1,  
275 stroke rate 2 and total stroke count were significantly greater in HEAT compared to CON ( $P < 0.05$ ,  $P <$   
276  $0.01$ ,  $P < 0.01$  respectively) (Table 1). Male participants showed a 1.01% improvement in the 50m  
277 performance in HEAT over CON ( $P < 0.05$ ); and stroke rate 1, stroke rate 2 and stroke count were  
278 higher in HEAT compared to CON ( $P < 0.01$ ,  $P < 0.05$ ,  $P < 0.01$ , respectively) (Table 1). For female  
279 participants, the 50m Freestyle times showed a trend to be 0.38% ( $P = 0.09$ ) faster in HEAT over CON,  
280 just under 0.4%- the value of the smallest worthwhile enhancement (13)- and stroke rate 2 and stroke  
281 count were higher ( $P < 0.05$ ,  $P < 0.1$  respectively) in HEAT compared to CON (Table 1). When the

282 50m Freestyle time was analyzed according to Hopkins (20), the practical inference of HEAT was  
283 'likely beneficial' (93.1%) for both genders combined and 'very likely beneficial' (97.5%) when  
284 looking at male participants alone. Female participants alone demonstrated a 'possible benefit' from  
285 HEAT, with any harmful negative effect from the condition being 'very unlikely' (0.9%).

286

### 287 3.3 PLYOMETRIC PRESS-UP

288 Absolute data for plyometric press-ups are shown in Fig. 3. Starting Strength and Peak Force were  
289 greater in HEAT compared to that in CON (Fig. 3) by 10.1% ( $P < 0.05$ ) and 10.7% ( $P = 0.097$ ).  
290 However, there was no difference in Peak Concentric Power when looking at all participants together  
291 (Table 1, Fig. 3). Male participants alone showed a 16.5%, 18.1% and 16.2% improvement in SS, PF  
292 and PCP, respectively, in HEAT over CON (all  $P < 0.01$ ). There was no difference found in female  
293 participant SS ( $P = 0.157$ ) or PF ( $P = 0.112$ ), though there was a trend in PCP ( $P = 0.07$ ) (Table 1).

294

### 295 3.4 TYMPANIC TEMPERATURE, SKIN TEMPERATURE & MUSCLE TEMPERATURE

296 There was no difference between conditions in mean torso  $T_{sk}$  before the warm-up or following the  
297 warm-up. After completion of the warm-up,  $T_{sk}$  had declined by  $\sim 4^{\circ}\text{C}$  in both conditions, with a slightly  
298 higher torso  $T_{sk}$  observed in CON compared to HEAT ( $29.5 \pm 1.1$  vs.  $29.1 \pm 1.0^{\circ}\text{C}$ ) ( $P < 0.05$ ).  
299 Following the recovery period, however,  $T_{sk}$  was  $2.3^{\circ}\text{C}$  higher in HEAT than CON ( $P < 0.001$ ). There  
300 was no difference in tympanic temperature between conditions (Table 2).  $T_m$  was estimated to be  
301  $36.7^{\circ}\text{C}$  in the HEAT condition in comparison to that of  $34.3^{\circ}\text{C}$  in the CON condition following the  
302 recovery period.

303

### 304 3.5 HR, RPE, THERMAL COMFORT AND SENSATION

305 There was no effect between conditions on either HR (Table 2) during the trials or RPE (17.5) for  
306 swimming performance. Thermal sensation was higher (towards 'hot') for HEAT compared to that for

307 CON between 5 minutes into the exercise and the end of the recovery period ( $P < 0.01$ ). Despite this  
308 difference, thermal discomfort for the conditions did not differ at any time points when observing all  
309 participants combined. There were no differences in thermal sensation between female and male  
310 participants ( $5.5 \pm 0.9$  vs  $5.4 \pm 0.6$ , respectively). There were no differences found in thermal  
311 discomfort scores between genders at the baseline, warm-up or post-50m Freestyle. However, a trend  
312 was observed in the magnitude of thermal discomfort scores between genders, with female participants  
313 showing a trend of scoring higher at 10- and 25-minutes during the HEAT recovery period ( $P = 0.056$ ,  
314  $P = 0.082$ , respectively).

#### 315 4. DISCUSSION

316 This study compared 50m freestyle performances and plyometric press-up measurements of a mixed-  
317 gender elite swimming group wearing heated jackets versus standard jackets for 30 minutes between the  
318 warm-up and racing, i.e. the period during competition events in which the swimmers tend to be in a  
319 holding area with limited ability to perform exercise. A trend for a relevant ( $>0.4\%$ ) magnitude (0.83%)  
320 of a 'likely beneficial' (300) improvement in the 50m Freestyle performance and a significantly  
321 improved force production was observed in the heated condition. Considering gender, the heated  
322 garment significantly improved the swimming performance of male participants by 1.01%, the effect  
323 being 'very likely beneficial', and also improving plyometric press-up measurements of both force and  
324 power production. However, the results for the female participants were less clear due to mixed results  
325 and the low number of participants in this group, showing a 'possible benefit' and 'unlikely to have  
326 harmful negative effects for the 50m times.

327

#### 328 4.1 Relevance to swimmers' routines

329

330 This study addresses the current issue raised by West *et al.* (7) of swimmers being unable to compete  
331 within the recommended timeframe of between 5- and 20-minutes after a warm-up, with time spent  
332 putting on the competition swimsuit, plus time in the holding areas often exceeding 30 minutes. Current  
333 literature examining the effects of different passive post warm-up procedures on swimming

334 performance is scarce and contradictory. Carlyle (31) demonstrated that swimmers who had an eight-  
335 minute hot shower or a ten-minute massage achieved 1% greater swim velocity than swimmers without  
336 any warm-up procedure; while conversely, De Vries (32) established that a ten-minute massage did not  
337 alter performance. McGowan *et al.*'s (15) recent study demonstrated that the passive heating (jacket)  
338 alone did not improve the overall 100m performance times (0.37%,  $P > 0.05$ ), though performance in  
339 the first half improved by 0.18%. However, the combination of passive heating as well as dryland-based  
340 activation exercises significantly improved time-trial performance ( $\sim 1.1\%$ ,  $P < 0.01$ ) (15). While  
341 Faulkner *et al.* (6, 12) and Raccuglia *et al.* (33) demonstrated for cyclists that heated tracksuit pants  
342 alone were sufficient in maintaining part of the warm-up  $T_m$  increase during a 30-minute transition  
343 period, or could even abolish the drop in  $T_m$  completely, the present study looked at testing the use of  
344 passive heating in swimmers with a more senior and elite group of participants, and for a shorter  
345 performance distance than in McGowan *et al.*'s (15) study. The positive effects observed in the present  
346 study are all around or above the 0.4% level of the smallest worthwhile enhancement in swimming (13),  
347 and the size of the improvements are similar to the combination strategy McGowan *et al.* tested. The  
348 observed results, especially for the male participants, give us great confidence in terms of practical  
349 relevance, as the analysis was a direct measure and simulation of a swimming race; and passive heating  
350 would therefore be recommended as a method to enhance a competition performance. Of the various  
351 differences between the present study and McGowan *et al.*'s study, the shorter distance of the event  
352 tested here (50m vs 100m) may be the most important, pointing to the impact of muscle heating mainly  
353 in short sprint type exercises, where central factors may be of little or no importance (34).

354  
355 The female participants demonstrated an average 0.38% ( $P = 0.09$ ) improvement in the 50m  
356 performance, just outside the set value of 0.4%. Though with only four data points the statistical power  
357 is low, the fact that half of the female participants improved in performance while half declined in  
358 performance indicates that even with a larger group, achieving a significant positive effect may be  
359 difficult, suggesting a potential gender effect.

360



361 As mean velocity is the product of the stroke rate and distance moved through the water with each  
362 completed stroke (Velocity [ $\text{m}\cdot\text{s}^{-1}$ ] = stroke rate [ $\text{s}^{-1}$ ] \* distance per stroke [m]), the increases observed  
363 in swimming velocity after wearing the heated garment are thought to be achieved mainly by the higher  
364 stroke rate (35). Greater stroke rates of 5% ( $P < 0.05$ ) and 3.8% ( $P < 0.01$ ) in stroke rate measures 1 and  
365 2 respectively were observed (Table 1). Studies have displayed that higher stroke rates have a clear  
366 relationship with an improved sprint freestyle performance (19, 36). An increase in stroke rate  
367 consequently decreased the distance per stroke, displayed in the 50m Freestyle by the higher stroke  
368 count in HEAT ( $P < 0.01$ ). The higher stroke rates are likely enabled by the greater preservation of  
369 muscle temperature between warm-up and performance under the HEAT condition.

370

## 371 4.2 MUSCLE TEMPERATURE

372

373 Although muscle temperature was not directly measured in the present study, it seems valid to suggest  
374 the HEAT strategy would have lessened the decline in muscle temperature following the completion of  
375 the warm-up (6, 12, 33). This is supported by the data which indicated that wearing the heated jacket  
376 following the warm-up for a 30-minute period raised  $T_{\text{sk}}$  by over  $2^{\circ}\text{C}$  more than  $T_{\text{sk}}$  without the heated  
377 jacket (Fig. 4). From this it is estimated that  $T_{\text{m}}$  was  $36.7^{\circ}\text{C}$  after HEAT in comparison to that of  $34.3^{\circ}\text{C}$   
378 in the CON condition (27), though the validity of De Ruiter *et al.*'s equation for the present application  
379 may be questioned. Nevertheless, based on the above, it is believed the post warm-up decline in  $T_{\text{m}}$  was  
380 smaller after the HEAT condition than that after the CON condition; and given that a difference in  $T_{\text{m}}$  as  
381 little as  $0.3^{\circ}\text{C}$  may critically affect performance (12), HEAT is assumed responsible for the positive  
382 effect on subsequent performances.

383

384 The majority of the beneficial effects of a warm-up have been attributed to temperature-related  
385 mechanisms (9). The relationship between muscle temperature and muscle function has been well  
386 established (10, 37, 11). Racinais & Oksa (5) concluded that muscle temperature may be the crucial  
387 factor in determining the outcome of short duration performance ( $R=0.91$ ). Therefore, maintaining a

388 raised muscle temperature through a warm-up is fundamental in achieving optimum sprint performance.  
389 Heightened muscle temperature enhances performance due to decreased stiffness of muscles and joints,  
390 increased transmission rate of nerve impulses, an altered force-velocity relationship and increased  
391 glycogenolysis, glycolysis and high-energy phosphate degradation (9). Thus, for a given force, the  
392 muscle-fiber conduction velocity should have increased following a heated recovery compared to that  
393 of the control (10). Greater muscle temperatures have also been linked to increases in myosin adenosine  
394 triphosphatase (ATPase) activity, increasing the rate of ATP turnover and calcium sequestration by the  
395 sarcoplasmic reticulum (5, 38). These physiological changes explain why a greater power output is  
396 achieved at higher muscles temperatures. As muscular power is a major factor in swimming success,  
397 determining the ability to generate propelling forces, it is vital that muscle temperature is maintained  
398 (39).

399  
400 Currently, there is no generally adopted method of maintaining muscle temperature during swimming  
401 competitions. Consequently, swimmers compete with less-than-optimal muscle temperatures, as warm-  
402 ups are generally completed from anywhere between 45 minutes to even 3 hours before racing. This is  
403 far from the optimum time frame of 5–20 minutes between cessation of warm-up and racing (2, 3); but  
404 due to lack of warm-up facilities and competition time constraints, optimizing the warm-up timing is  
405 not feasible. Durations longer than the suggested window to compete result in lower-than-optimal  
406 muscle temperatures, as postulated in the control condition. This will subsequently effect muscle  
407 contractile properties, producing slower, less powerful contractions (27, 37). As a result, swimmers may  
408 not present themselves at the optimum physical condition, thus decreasing their chances of achieving  
409 their greatest performance times. Any improvement that can be achieved would provide the individual  
410 swimmer with a competitive advantage.

411

#### 412 4.3 PLYOMETRIC PRESS-UP DATA

413

414 In order to analyze the arm forces and power production separately from other factors affecting  
415 swimming performance, plyometric press-ups were assessed. The focus of the study was based on  
416 upper-body measurements, as arm strength is the main criterion used to explain sprint swimming  
417 performance (16). The muscle activation required during a press-up involves three of the four main  
418 swimming muscles used to propel a swimmer through the water: pectoralis major, deltoids and triceps  
419 brachii (26). Hence, press-up measurements are assumed to be a valid indicator of swimming  
420 performance, requiring the same muscle groups as a bench press (except in the prone position), which  
421 has been associated with swimming velocity (16). This, along with the pilot study, displayed plyometric  
422 press-up reliability and validity as a functional measurement of upper-body power output within  
423 swimmers.

424

425 While the male participants showed clear and substantial improvements due to the use of the heated  
426 garment (>15%), the female participants did not display any improvements in peak concentric power or  
427 force production after HEAT relative to CON. This may be due to the physiological differences  
428 between the female and male participants, and the differences in weekly exercise routines (Time in  
429 gym: Male:  $5.5 \pm 0.4$  vs Female:  $2.6 \pm 1.1$  hours). But again, due to the small sample size of female  
430 participants, the chance of type II error is dramatically increased.

431

432 It can be assumed that the levels of greater force and power observed in the plyometric push-ups would  
433 have transferred into the 50m Freestyle performance. Whereby, improvements in arm strength may  
434 result in higher levels of maximum force per stroke and greater power would increase the stroke rate,  
435 producing faster swimming velocities displayed in the study (16). These findings are consistent with  
436 previous literature, presenting a positive relationship between the body temperature's effect on  
437 movement velocity and performance (5). This study supports previous literature in that, with every  $1^{\circ}\text{C}$   
438 rise in muscle temperature, there is an estimated 4%–10% improvement in peak power output; as in the  
439 present study,  $T_m$  is thought to be  $\sim 2^{\circ}\text{C}$  greater in HEAT than that of CON (10, 11, 12), with a 16% to

440 18% increase in SS, PF and PCP. The studies to date assessing passive heating have focused on  
441 assessing lower-body measurements of power output. To the authors' knowledge, this is the first study  
442 to assess passive heat maintenance on upper-body measurements of force production and power output;  
443 therefore, the enhancements observed are an important addition to the literature.

444

#### 445 **Thermal sensation and comfort**

446 It can be seen as positive that the only difference in subjective measurements was thermal sensation ( $P$   
447  $< 0.01$ ) between the two conditions, likely due to the increased  $T_{sk}$  ( $P < 0.0001$ ) (Table 2) after HEAT.  
448 However, importantly, thermal comfort (Table 2) did not differ, signifying that the participants regarded  
449 both the heating and the lack-of-heating conditions as thermally acceptable. The conformity of the  
450 perceived comfort is fundamental, as pre-exercise thermal discomfort has been associated with impaired  
451 performance (40). However, when comparing gender scores, female participants rated significantly  
452 higher for thermal discomfort over the 30-minute recovery period under the HEAT condition,  
453 suggesting they were slightly uncomfortable in comparison to the male participants showing ratings of  
454 discomfort ( $2.9 \pm 0.4$  vs  $1.6 \pm 0.3$ , HEAT vs CON, respectively  $P < 0.05$ ). This may have affected the  
455 female participants' performances, as only two out of the four female participants demonstrated  
456 improvements in the 50m Freestyle performance under the HEAT condition.

457

458 This variance may be due to morphology differences in body size and body composition between the  
459 genders affecting thermoregulation. As the female participants had a significantly higher body-surface-  
460 area to mass-ratio and body-fat percentage than the male participants, they may have experienced a  
461 greater heat strain (41). However, as tympanic temperature did not significantly differ between genders  
462 and both were far from  $39^{\circ}\text{C}$ , the higher ratings for thermal discomfort may not be due to heat strain  
463 (5). Instead, the female participants may have felt slightly more uncomfortable, possibly due to a higher  
464 thermoreceptor density based on having a significantly lower body-surface area (42). Females are more  
465 sensitive to innocuous heat ( $40^{\circ}\text{C}$ ) stimulation than males (42). Consequently, the differences observed

466 in performances between male and female participants may not be due to differences in the thermal  
467 state of the body, but to thermal perception. As performance intensity is strongly influenced by the  
468 thermal status of the body, detected by thermal comfort, this may have had a negative impact on the  
469 female performances (40). This highlights the importance of studies testing both male and female  
470 participants rather than generalizing findings of both genders. Consequently, females may favor a  
471 reduced heating power and reduced temperature for the heated jackets, a point for further study.

472

#### 473 4.4 LIMITATIONS

474

475 Measures of  $T_m$  were not recorded during the trials due to its invasive nature and the problems with  
476 keeping sterility when entering the pool. This does not detract from the meaningfulness of this data, as  
477 although attenuation of the  $T_m$  drop is vital for performance enhancement, estimates of  $T_m$  were  
478 calculated based on its linear relationship with  $T_{sk}$  (27, 43). Also, given that previous studies from our  
479 lab, most recently Faulkner *et al.* (6, 12) and Raccuglia *et al.* (33), have demonstrated  $T_m$  maintenance  
480 with the use of passive external heating, it is highly likely that  $T_m$  in the present study would have  
481 followed similar time course changes. In addition, another limitation to this study was a relatively small  
482 number of female subjects tested ( $n=4$ ) due to logistical problems, increasing the risk of type II errors  
483 and limiting the possibility of gender comparisons. To confirm the observed response differences  
484 between genders, future research should test a greater number of female swimmers, with the possibility  
485 of a self-adjustable temperature control in order to avoid any possible negative effects of thermal  
486 discomfort and subsequent negative impacts on performance.

487

#### 488 4.5 CONCLUSIONS

489

490 This study has demonstrated that a 30-minute period of upper-body external heating using electrically  
491 heated jackets post warm-up leads to a significant and relevant improvement in sprint-swimming  
492 performance, upper-body force and power output when compared to a non-heated control in elite male

493 swimmers. No significant effect was observed for the female group on its own, suggesting a gender  
494 difference with possible links to gender differences in experienced discomfort; but given the small  
495 female group size, further research should be carried out. This study provides an important practical  
496 application of heated garments for swimmers due to the unavoidable timeframe between completion of  
497 warm-up and racing. These findings may be relevant to all sports that experience delays after warm-up  
498 or have an intermittent nature, and are reliant on high peak-power output, as in sprint exercises. Given  
499 that the jackets used in this testing were mainly designed for thermal comfort in motorcyclists, it is  
500 hypothesized that the heating provided can be further improved by designing the jackets' heater  
501 distribution to align with the major muscle groups used in swimming. Additional leg heating (though  
502 limited impact is expected in freestyle) could also be considered, e.g. for breaststroke. In addition,  
503 personal control of the heating power may contribute positively to the acceptance of the heating jackets,  
504 especially for female participants.

505

## 506 5 PRACTICAL IMPLICATIONS

507 This study supports the use of heated garments for the upper body to be used by competitive swimmers  
508 to maintain muscle temperature between the warm-up and the event, or between events, in order to  
509 improve performance. More work is needed to understand why this benefit was evident in male  
510 participants but not clear in the tested female participants. It is possible that females' higher sensitivity  
511 to heat could influence the benefit of the used warming procedure, and would require an adjustable  
512 heating system.

## 513 ACKNOWLEDGMENTS

514 The authors acknowledge the support by Speedo in the procurement of the heating garments and by  
515 British Swimming for use of relevant measuring devices. Furthermore, all swimmers who participated  
516 in the study and volunteers who helped with testing are thanked for their contribution.

## 517 CONFLICT OF INTEREST

518 No conflicts of interest are reported.

## 519 6. REFERENCES

520

- 521 1. Fradkin, AJ, Zazryn, TR, Smoliga, JM. Effects of warming-up on physical performance: a  
522 systematic review with meta-analysis. *J Strength Cond Res.* 2010; 24(1): 140–148.
- 523 2. Zochowski, T, Johnson, E, Sleivert, GG. Effects of varying post-warm-up recovery time on  
524 200-m time-trial swim performance. *Int J Sports Physiol Perform.* 2007; 2: 201–211.
- 525 3. Bishop, D. Potential Mechanisms and the Effects of Passive Warm-up on Exercise  
526 Performance. *Sports Med.* 2003; 33(6): 439–454.
- 527 4. Russell M, West DJ, Briggs MA, Bracken RM, Cook CJ, Giroud T, Gill N, Kilduff LP. A  
528 passive heat maintenance strategy implemented during a simulated half-time improves lower  
529 body power output and repeated sprint ability in professional Rugby Union players. *PLoS One.*  
530 2015 Mar 18;10(3):e0119374. doi: 10.1371/journal.pone.0119374. eCollection 2015.
- 531 5. Racinais S, Oksa J. Temperature and neuromuscular function. *Scand J Med Sci Sports.* 2010;  
532 20: 1–18.
- 533 6. Faulkner, SH, Ferguson, RA, Gerrett, N, Hupperets, M, Hodder, SG, Havenith, G. Reducing  
534 muscle temperature drop after warm-up improves sprint cycling performance. *Med Sci Sports  
535 Exerc.* 2013; 45(2): 359–365.
- 536 7. West, DJ, Dietzig, BM, Bracken, RM, Cunningham, DJ, Crewther, BT, Cook, CJ. Influence of  
537 post-warm-up recovery time on swim performance in international swimmers. *J Sci Med Sport.*  
538 2013; 16(2): 172–176.
- 539 8. Mohr, M, Krstrup, P, Nybo, L, Nielsen, JJ, Bangsbo, J. Muscle temperature and sprint  
540 performance during soccer matches - Beneficial effect of re-warm-up at half-time. *Scand J Med  
541 Sci Sports.* 2004; 14(i): 156–162.
- 542 9. Bishop, D. Warm-up II Performance Changes Following Active Warm-up and How to  
543 Structure the Warm-up. *Sports Med.* 2003; 33(7): 483–498.
- 544 10. Bergh, U, Ekblom, B. Influence of muscle temperature on maximal muscle strength and power

- 545 output in human skeletal muscles. *Acta Physiol Scand.* 1979; 107(1): 33–37.
- 546 11. Sargeant, AJ. Effect of muscle temperature on leg extension force and short-term power output  
547 in humans. *Journal of Applied Physiology and Occupational Physiology.* 1987; 56(6): 693–698.
- 548 12. Faulkner, SH, Ferguson, RA, Hodder, SG, Havenith, G. External muscle heating during warm-  
549 up does not provide added performance benefit above external heating in the recovery period  
550 alone. *Eur J Appl Physiol.* 2013; 113(11): 2713–2721.
- 551 13. Pyne, D, Trewin, C, Hopkins, W. Progression and variability of competitive performance of  
552 Olympic swimmers. *J Sports Sci.* 2004; 22: 613–620.
- 553 14. Deschodt, JV, Arzac, LM, Rouard, AH. Relative contribution of arms and legs in humans to  
554 propulsion in 25-m sprint front-crawl swimming. *Eur J Appl Physiol.* 1999; 80(3): 192–199.
- 555 15. McGowan, CJ, Thompson, KG, Pyne, DB, Raglin, JS, Rattray, B. Heated jackets and dryland-  
556 based activation exercises used as additional warm-ups during transition enhance sprint  
557 swimming performance. *J Sci Med Sport.* 2015; Online First. doi:10.1016/j.jsams.2015.04.012.
- 558 16. Morouço, PG, Marinho, DA, Marques, MC. Are dry-land strength metrics and forces exerted  
559 in-water related with high swimming velocity in young athletes? *International Society of*  
560 *Biomechanics in Sport.* 2014; 32 International Conference of Biomechanics in Sports, Johnson  
561 City, US
- 562 17. Carl, DL, Leslie, N., Dickerson, T., Griffin, B., Marksteiner, A. Bench Press and Leg Press  
563 Strength and its Relationship with In-Water Force and Swimming Performance when Measured  
564 in-season in Male and Female Age-group Swimmers, in: XI International Symposium 2010;  
565 [http://www.nih.no/upload/BMS2010/Documents/BMS2010\\_BMS\\_XI\\_final\\_lowres.pdf](http://www.nih.no/upload/BMS2010/Documents/BMS2010_BMS_XI_final_lowres.pdf)
- 566 18. Cossor JM, Blanksby BA, Elliott BC. The influence of plyometric training on the freestyle  
567 tumble turn. *J Sci Med Sport.* 1999 Jun;2(2):106-16.
- 568 19. Costa, MJ, Bragada, JA, Marinho, DA, Silva, AJ, Barbosa, TM. Longitudinal interventions in  
569 elite swimming: a systematic review based on energetics, biomechanics, and performance. *J*  
570 *Strength Cond Res.* 2012; 26(7): 2006–2016.
- 571 20. Hopkins, WG. Spreadsheets for analysis of controlled trials with adjustment for a predictor.  
572 *Sportscience.* 2007; 10: 1-5.



- 573 21. Ericsson, KA, Lehmann, AC. Expert and exceptional performance: evidence of maximal  
574 adaptation to task constraints. *Annu Rev Psychol.* 1996; 47: 273–305.
- 575 22. Hogarth, L, Deakin, G, Sinclair, W. Are plyometric push-ups a reliable power assessment tool?  
576 *Journal of Australian Strength and Conditioning.* 2013; 21(1): 67–69.
- 577 23. Cronin, JB, Hing, RD, McNair, PJ. Reliability and validity of a linear position transducer for  
578 measuring jump performance. *J Strength Cond Res.* 2004; 18(3): 590–593.
- 579 24. ASHRAE (American Society for Heating Refrigeration and Air-conditioning Engineers.  
580 Chapter 9: Thermal Comfort. In: *ASHRAE Handbook of Fundamentals.* Atlanta (GA),  
581 ASHRAE. 9.1-9.32, 2013.
- 582 25. Havenith, G. Laboratory assessment of cold weather clothing, in: *Textiles for cold weather*  
583 *apparel*, Edited by J. Williams, Woodhead Publishing in Textiles: Number 93; 2009; Woodhead  
584 Publishing Limited, Oxford Cambridge New Delhi
- 585 26. Birrer, P. The Shoulder, EMS and the Swimming Stroke. *The Journal of Swimming Research.*  
586 1986; 2(1): 20–23.
- 587 27. De Ruiter, CJ, Jones, AD, Sargeant, AJ, De Haan, A. Temperature effect on the rates of  
588 isometric force development and relaxation in the fresh and fatigued human adductor pollicis  
589 muscle. *Exp Physiol.* 1999; 84: 1137–1150.
- 590 28. Wurm, B, VanderZanden, T, Spadavecchia, M et al. Kinetic analysis of several variations of  
591 push-ups. *J Strength Cond Res.* 2010; 25(10): 2891-2894.
- 592 29. Borg, GA. Psychological bases of perceived exertion. *Med Sci Sports Exerc.* 2001; 14: 377–  
593 381.
- 594 30. Hopkins, WG, Marshall, SW, Batterham, AM, Hanin, J. Progressive statistics for studies in  
595 sports medicine and exercise science. *Med Sci Sports Exer.* 2009; 41(1): 3–13.
- 596 31. Carlyle, F. Effect of preliminary passive warming on swimming performance. *Res Q Exerc*  
597 *Sport.* 1956; 27(2): 143–151.
- 598 32. De Vries, HA. Effects of various warm-up procedures on 100-yard times of competitive  
599 swimmers. *Res Q.* 1959; 30: 11–22.
- 600 33. Raccuglia, M, Lloyd, A, Filingeri, D, Faulkner, S, Hodder, S, Havenith, G. Post-warm-up

- 601 muscle temperature maintenance: blood flow contribution and external heating optimization,  
602 European Journal of Applied Physiology, February 2016, Volume 116, Issue 2, pp 395-404  
603 ISSN: 1439-6327.
- 604 34. Lloyd A., Hodder S., Havenith G. The interaction between peripheral and central fatigue at  
605 different muscle temperatures during sustained isometric contractions. *Am J Physiol Regul*  
606 *Integr Comp Physiol.* 2015 Jun 3;ajpregu.00061.2015. doi: 10.1152/ajpregu.00061.2015. 2015  
607 Aug 15;309(4):R410-420.
- 608 35. Craig, AJ, Pendergast, D. Relationships of stroke rate, distance per stroke, and velocity in  
609 competitive swimming. *Med Sci Sports Exerc.* 1979; 11(3): 278–283.
- 610 36. Toussaint, HM, Beek, PJ. Biomechanics of competitive front crawl swimming. *Sports Med.*  
611 1992; 13(I):8–24.
- 612 37. Davies, CT, Mecrow, IK, White, MJ. Contractile properties of the human triceps surae with  
613 some observations on the effects of temperature and exercise. *Eur J Appl Physiol.* 1982; 49(2):  
614 255–269.
- 615 38. Stein, RB, Gordon, T, Shriver, J. Temperature dependence of mammalian muscle contractions  
616 and ATPase activities. *Biophys J.* 1982; 40(2): 97–107.
- 617 39. Sharp, RL, Troup, JP, Costill, DL. Relationship between power and sprint freestyle swimming.  
618 *Med Sci Sports Exerc.* 1982; 14(1): 53–56.
- 619 40. Schlader, ZJ, Simmons, SE, Stannard, SR, Mündel, T. Skin temperature as a thermal controller  
620 of exercise intensity. *Eur J Appl Physiol.* 2011; 111(8): 1631–1639.
- 621 41. Havenith, G. Human surface to mass ratio and body core temperature in exercise heat stress - A  
622 concept revisited. *Journal of Thermal Biology.* 2001; 26(4-5): 387–393.
- 623 42. Gerrett, N, Ouzzahra, Y, Coleby, S et al. Thermal sensitivity to warmth during rest and  
624 exercise: A sex comparison. *Eur J Appl Physiol.* 2014 114: 1451–1462.
- 625 43. De Ruiter, CJ, De Haan, A. Temperature effect on the force/velocity relationship of the fresh  
626 and fatigued human adductor pollicis muscle. *Eur J Physiol.* 2000; 440(1): 163–170.
- 627

628 **Table Captions:**

629 Table 1. Male and Female (n=12), Male only (n=8), Female only (n=4), 25m split, 50m Time, Stroke  
630 Rate 1 (SR1), Stroke Rate 2 (SR2), Stroke Count (SC).

631

632 Table 2. Tympanic temperature (TT), skin temperature ( $T_{sk}$ ), HR, thermal sensation (TS) and thermal  
633 comfort (TC) at baseline (BASE), after warm-up (30 WUP), after passive recovery (30REC) and  
634 straight after the maximal 50m Freestyle (POST50) for control (CON) and heating (HEAT) conditions  
635 (n=12).

636

637 **Figure Captions**

638 Figure 1. Thermogram of Powerlet rapidFIRE Proform Heated Jacket Liner

639

640 Figure 2. Mean and Individual 50m Freestyle swimming performance times for control (CON) and  
641 heating (HEAT) for Males and Females. \*  $P < 0.05$ , HEAT < CON. §  $P < 0.1$ , HEAT < CON.

642

643 Figure 3. Mean ( $\pm$ SD) values of starting strength (SS) and peak force (PF) and Peak Concentric Power  
644 (PCP) for control (CON) and heating (HEAT). §  $P < 0.1$  HEAT > CON. \* $P < 0.05$ , HEAT > CON. \*\*  
645  $P < 0.01$ , HEAT > CON.

646

647 Figure 4. Mean (SD) upper-body mean skin temperature measures prior to warm-up (0WUP), straight  
648 after warm-up (30WUP), and after 30 minutes seated recovery (30REC) in control (CON) and heating  
649 (HEAT) (n=12). \*  $P < 0.05$ , CON > HEAT. †  $P < 0.0001$ , HEAT > CON.

650

651