

Loughborough University
Institutional Repository

*Asymmetric total synthesis
of (-)-erogorgiaene and its
C11 epimer and
investigation of their
antimycobacterial activity*

This item was submitted to Loughborough University's Institutional Repository by the/an author.

Citation: INCERTI-PRADILLOS, C.A. ... et al, 2016. Asymmetric total synthesis of (-)-erogorgiaene and its C11 epimer and investigation of their antimycobacterial activity. *Chemistry - A European Journal*, 22 (40), pp. 14390-14396.

Additional Information:

- This is the peer reviewed version of the following article: INCERTI-PRADILLOS, C.A. ... et al, 2016. Asymmetric total synthesis of (-)-erogorgiaene and its C11 epimer and investigation of their antimycobacterial activity. *Chemistry - A European Journal*, 22 (40), pp. 14390-14396., which has been published in final form at <http://dx.doi.org/10.1002/chem.201602440>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

Metadata Record: <https://dspace.lboro.ac.uk/2134/22370>

Version: Accepted for publication

Publisher: © Wiley

Rights: This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Please cite the published version.

Asymmetric Total Synthesis of (–)-Erogorgiaene and Its C11 Epimer and Investigation of Their Antimycobacterial Activity

Celia A. Incerti-Pradillos^[a], Mikhail A. Kabeshov^[a], Paul S. O'Hora^[a], Sergei A. Shipilovskikh^[a,b], Aleksandr E. Rubtsov^[b], Vera A. Drobkova^[b], Svetlana Yu. Balandina^[b], and Andrei V. Malkov^{[a]*}

Abstract: A short, 9-step, highly enantioselective synthesis of (–)-erogorgiaene and its C11 epimer is reported. The key stereochemistry controlling steps involve catalytic asymmetric crotylation, anionic oxy-Cope rearrangement and cationic cyclization. (–)-Erogorgiaene exhibited promising antitubercular activity against multidrug resistant strains of *M. tuberculosis*.

Introduction

For many years, natural products served as an inspirational source for developing new drugs. Now, with antibiotics losing their effectiveness at an alarming rate, there is an urgent need to find new antibacterial leads to tackle this problem. Diterpenoids bearing a serrulatane (biflorane) skeleton exhibited a wide range of useful biological properties including promising antimycobacterial activity.^[1] Selected representatives of serrulatane metabolites found in the marine gorgonian *Pseudopterogorgia elisabethae* (**1–4**) and in terrestrial plants, e.g. Mexican evergreen shrub *Leucophyllum frutescens* (**5**) and Australian *Eremophila* (**6**), are shown in Figure 1. Inspection of the stereochemical elements of these compounds reveals that the diterpenes isolated from the sea organisms differ from the terrestrial counterparts by the relative configuration at C-11. Additionally, these two groups of serrulatane diterpenes generally belong to the opposite enantiomeric series, although the levorotatory enantiomer of **1** was isolated from a brown alga (*Dictyota dichotoma*).^[2] The relatively simple structures and potent biological activity of serrulatane diterpenes that in many instances is augmented by the low abundance and scarcity of supply from their natural sources make them attractive synthetic targets for exploring their therapeutic potential. However, the lack of functional groups near the stereogenic centres in the predominantly hydrocarbon serrulatane core represents a significant challenge for stereose-

lective synthesis. The successful strategies for controlling stereochemistry in the total synthesis of this family of diterpenes include chiral pool,^[3] chiral reagents/chiral substrates,^[4] and asymmetric catalysis.^[5] In the case of erogorgiaene **1**, the synthesis usually required 11 steps or more, for a notable exception of the 8-step convergent sequence developed by Aggarwal,^[4b] which also provided access to other three diastereoisomers of **1**. Recently, we reported on enantioselective synthesis of (–)-elisabethadione **3**, where the key stereochemistry defining steps included (i) catalytic asymmetric crotylation, (ii) anionic oxy-Cope rearrangement (AOC) and (iii) cationic cyclization.^[5d] Herein, we present extension of this strategy to a wider range of serrulatane diterpenes with a detailed insight into the stereochemistry of the AOC. The generality of the method is illustrated by a short, highly enantioselective synthesis of both C-11 diastereoisomers of erogorgiaene **1** followed by investigation into their activity against *Mycobacterium tuberculosis* including multidrug resistant strains.




Figure 1. Serrulatane metabolites.

Results and Discussion


Retrosynthetic analysis of erogorgiaene **1** (Scheme 1) identifies α -methylcinnamaldehyde **11b** as the starting material, which

[a] Dr. C. A. Incerti-Pradillos, Dr. M. A. Kabeshov, Dr. P. S. O'Hora, S. A. Shipilovskikh, and Prof. Dr. A. V. Malkov
Department of Chemistry
Loughborough University
Loughborough, LE11 3TU, UK
Fax: +44 (0)1509 22 3925
E-mail: A. Malkov@lboro.ac.uk

[b] S. A. Shipilovskikh, Dr. V. A. Drobkova, S. Yu. Balandina and Dr. A. E. Rubtsov
Department of Chemistry
Perm State University
Bukireva 15, Perm, 614990, Russia


Supporting information for this article is given via a link at the end of the document

undergoes asymmetric allylation with **10** to afford alcohol **9** followed by the anionic oxy-Cope rearrangement into **8** that installs two out of three stereogenic centers of the serrulatane skeleton. Ensuing cationic cyclization after side chain extension delivers the final stereogenic center (**7**).




Scheme 1. Retrosynthetic analysis of erogorgiaene **1**.

Catalytic methods for highly enantio- and diastereoselective crotylation of unsaturated aldehydes mediated by chiral Lewis bases have been reported from this laboratory and by others.^[5d, 6] Therefore, the initial studies for assessing feasibility of this sequence were focused on the stereochemical course of the AOC rearrangement and subsequent transformations. The model experiments were carried out with alcohols *syn-9a* and *anti-14* that are conveniently synthesized in diastereomerically pure form by crotylation of α -methylcinnamaldehyde **11a** with *Z*-crotyltrichlorosilane **10** and its *E*-isomer **13**, respectively^[5d, 6] (Scheme 2). The enolate resulting from the rearrangement was protonated with MeOH at -78°C and the intermediate aldehyde, without isolation, was subjected to Wittig alkenylation to furnish **15a**. Clearly, *syn-9a* exhibited much better stereoselectivity. Note that the AOC is responsible for the formation of C4 and the geometry of the alkene (*E/Z*) in **15a**, whereas C11 is formed post rearrangement, in the protonation of the enolate. The charge-accelerated [3,3] sigmatropic process is likely to proceed via a chair-like transition state, however chirality transfer in the AOC is not straightforward and strongly depends on the substitution pattern of the substrate.^[7] Analysis of the stereochemical course of the AOC is presented in Scheme 3.



Scheme 2. Model studies of anionic oxy-Cope rearrangement.

In the rearrangement of *syn-9a*, the two possible transition states **A** and **B** will result in the respective products **17** and **18** with a different alkene geometry and, importantly, with the opposite absolute configuration at the benzylic centre. However, the TS **B** is clearly disfavored due to the axial arrangement of the three adjacent substituents: the two methyl groups and the alkoxide. Therefore, enolate **17** is formed preferentially (**17/18** 25:1). Both chair-like transition states **C** and **D** resulting from *anti-14* feature two axial and one equatorial substituent, however TS **D** appears to be much less favourable due to the adverse 1,3-diaxial interactions between the methyl groups. The observed lack of stereoselectivity in this instance (**17'/18'** 3:1) suggests that a boat TS might be involved.^[7a] It is also pertinent to note that to produce the same C-4 enantiomer of **17/17'**, alcohols *syn-9a* and *anti-14* should come from the opposite enantiomeric series (Schemes 2 and 3).




Scheme 3. Stereochemistry of anionic oxy-Cope rearrangement.

As a consequence, to enable an efficient transfer of chirality in the asymmetric variant of the AOC, it is a prerequisite to employ diastereomerically pure **9a** since asymmetric crotylation is usually face-selective and affords *syn* and *anti* isomers (**9** and **14**) with the same absolute configuration of the alcohol.^[5d]

Furthermore, *E* configuration of the double bond in **17** (and **15a**) is crucial for achieving 1,4-*trans* tetralin structure **16a** in the subsequent cationic cyclization (Scheme 2, cf Figure 1), as *Z*-alkene **18** appears to favour the respective 1,4-*cis* tetralin.^[5d, 8]

While the engineered substitution pattern in **9a** ensures the high stereochemical fidelity in the formation of the C-4 stereogenic center during the AOC, formation of the C-11 relies on a face-selective protonation of the potassium enolate resulting from the rearrangement. Methanol quenching at $-78\text{ }^{\circ}\text{C}$ furnishes a nearly equimolar mixture of isomeric *syn*-**19** and *anti*-**20**. Preliminary calculations at the semi-empirical level^[9] identified two major conformations **E** and **F** (Scheme 4) of the intermediate enolate (modelled as enol). A less than 0.3 kcal/mol energy difference between **E** and **F** correlates with the observed lack of stereoselectivity in the protonation. A brief optimisation of quenching conditions under kinetic control revealed that protonation of the enolate with a bulkier 2,4,6-*tert*-butylphenol at $-78\text{ }^{\circ}\text{C}$ gave an improved 2.5:1 d.r. (Scheme 2). It is worth noting that equilibration of a 2:1 **19/20** mixture with MeONa in MeOH/THF at rt for 12 h resulted in a 1:1.3 mixture favoring **20**.




Scheme 4. Diastereoselectivity of protonation of enolate **17**.

Next, we attempted a racemic variant of the synthesis of erogorgiaene **1** starting from α -methylcinnamaldehyde **11b** (Scheme 5). Crotylation of **11b** with *Z*-**10** using DMF as a Lewis base activator furnished alcohol *syn*-**9a** in 64% yield, which was then subjected to the AOC. Methanol quenching followed by Wittig chain extension afforded methyl ester **15b** as a 1:1 mixture of C11 diastereoisomers in 30% yield (unoptimised). Treatment of **15b** with methanesulfonic acid in DCM at $-78\text{ }^{\circ}\text{C}$ followed by a slow warming to room temperature yielded an inseparable mixture of *syn/anti* and regioisomers **16b/21** in nearly equimolar quantities making this route impractical. Note that the problem of controlling regioselectivity in the cationic cyclization did not exist in the previously reported synthesis of elisabethadione **3** since only one position on the aromatic ring was available.^[5d]

Therefore, an appropriate protection strategy was considered next. In a revised route (Scheme 6), 2-bromo substituted cinnamaldehyde **11c** was chosen as the substrate, the bromine

serves as a blocking group to eliminate the issue of regioselectivity in the cationic cyclization. Bromine has a successful record of being employed as a protecting group in total synthesis of natural products on a large scale;^[10] it can be readily removed by reduction once it has served its purpose. Provided the debromination could be combined with other reduction process, the revised strategy would not affect the overall number of steps in the synthesis.



Scheme 5. Model studies towards the synthesis of (±)-erogorgiaene (**1**).

The route commenced with the synthesis of **11c** by aldol condensation of commercially available aldehyde **22** with propional (70%). Asymmetric crotylation of **11c** with **10** catalysed by our recently developed bis-*N*-oxide (–)-**12**^[5d] (2 mol%) in propionitrile at $-60\text{ }^{\circ}\text{C}$ on a 5 mmol scale furnished homoallylic alcohol *syn*-(–)-**9c** in 72% yield and excellent 99:1 er (d.r. > 25:1). The AOC rearrangement was carried out following the conditions described for alcohol **9a**, except the reaction was performed in DME at RT. Quenching the resulting enolate with 2,4,6-tri-*tert*-butylphenol at $-78\text{ }^{\circ}\text{C}$ gave a 2:1 ratio of C-11 diastereoisomers. To minimise manipulation of the oxidation-prone aldehyde and to avoid epimerisation at C-11, the crude aldehyde was subjected to Horner-Wadsworth-Emmons alkenylation conditions^[11] to furnish unsaturated ester **15c** (81% over 2 steps). The diastereoisomers were not separable at this point therefore the synthesis continued with the mixture. Ester **15c** was submitted to the cationic cyclisation with MeSO₃H to produce *trans*-tetraline **16c** in 83% yield and good stereoselectivity (dr 14:1). With all the stereocentres in place, the next task was to reduce the unsaturated ester to the saturated alcohol and remove Br from the aromatic ring. After some experimentation it was found that continuous flow hydrogenation in methanol using H-cube (2.0 mL/min, 20 bar, 40 $^{\circ}\text{C}$, 4 h) equipped with a 10% Pd/C catalyst cartridge accomplished both the double bond reduction and debromination. The solvent was replaced for DCM and the

saturated ester was reduced with DIBAL-H into primary alcohol **7** in 70% overall yield. Preparative HPLC separation of the diastereoisomers furnished pure (–)-**7** and its C-11 epimer (–)-**7a**. Dess-Martin oxidation of (–)-**7** led to the respective aldehyde in 84% yield. Finally, Wittig alkenylation afforded (–)-erogorgiaene **1** (dr 12:1) in 80% yield exhibiting optical rotation $[\alpha]_D = -25.5$ (c 0.52, CHCl_3) that is in agreement with the value for the natural product isolated from a brown alga (*Dictyota dichotoma*): $[\alpha]_D = -28.0$ (c 0.05, CHCl_3)^[12] and the absolute value of the dextrorotatory enantiomer from *Pseudopterogorgia elisabethae*: $[\alpha]_D = +24.4$ (c 3.2, CHCl_3).^[1a] Synthesis of the C-11 epimer (–)-**1a** (dr 7:1) from the respective alcohol (–)-**7a** was completed through the same two-step sequence.


According to an earlier report,^[1a] (+)-erogorgiaene **1** exhibited promising activity against *Mycobacterium tuberculosis* H₃₇Rv (96% growth inhibition at 12.5 $\mu\text{g/mL}$). Due to the minute quantities of the isolated material, it has not been further investigated. C-11 epimer (–)-**1a** has the same relative and absolute configuration as (–)-leubethanol **5**, which showed a potent activity against both drug-sensitive and multidrug resistant (MDR) strains at minimal inhibitory concentrations (MIC) in the range of 6.25–12.5 $\mu\text{g/mL}$.^[12] Therefore, with the two diastereoisomers (–)-**1** and (–)-**1a** in hand, we next evaluated their antitubercular activity against two clinical isolates of *M. tuberculosis*, one of

which showed sensitivity to isoniazid and rifampicin, the first-line anti-TB drugs, the other exhibited multidrug resistance. Selected data are presented in Table 1.

Table 1. Antimycobacterial activity of (–)-**1** and (–)-**1a**

| Entry | Compound | MIC ($\mu\text{g/mL}$) | |
|-------|----------------|--------------------------|------------|
| | | Susceptible to TB drugs | MDR Strain |
| 1 | (–)- 1 | 31,25 | 125,0 |
| 2 | (–)- 1a | 62,5 | 250,0 |

The results revealed that (–)-**1** was twice as active as its C-11 epimer (–)-**1a** against both TB strains. Comparison with the literature data for erogorgiaene **1** indicates that the dextrorotatory enantiomer (+)-**1** isolated from the octocoral *Pseudopterogorgia elisabethae* showed somewhat higher activity than its enantiomer (–)-**1** synthesised in this work. High potency of (–)-leubethanol **5** seems to indicate the importance of the hydroxyl group in the position 8 of the aromatic ring for antitubercular activity. It is worth noting that the both C-11 epimers induced growth inhibition of the MDR strain and can serve as leads for further development.



Scheme 6. Asymmetric total synthesis of (–)-**1** and its C-11 epimer (–)-**1a** (ArOH = 2,4,6-(*t*-Bu)₃C₆H₂OH).

Conclusions

A short, 9-step synthesis of natural (–)-erogorgiaene (10.5% overall yield) and its C-11 epimer from a commercially available precursor has been accomplished through a sequence of simple transformations. The key reactions for introducing stereogenic centers include (i) highly enantioselective catalytic crotylation, (ii) anionic oxy-Cope rearrangement and (iii) cationic cyclisation. In the latter instance, protection strategy was seamlessly integrated into the synthesis without altering the number of steps. The compounds showed promising activity against clinical isolates of *M. tuberculosis* including multidrug resistant strain. Ready access to α -methyl cinnamaldehydes with various substitution patterns in the aromatic ring opens the door to the synthesis of natural and unnatural analogues of these diterpenes for biological evaluation.

Experimental Section

General Methods and Materials.

The NMR spectra were recorded in CDCl₃, ¹H at 400 MHz and ¹³C at 100.6 MHz with chloroform-d₇ (δ 7.26, ¹H; δ 77.0, ¹³C) as internal standard unless otherwise indicated. Various 2D-techniques and DEPT experiments were used to establish the structures and to assign the signals. The IR spectra were recorded in KBr disc unless otherwise indicated. The mass spectra were recorded in a positive or negative ion mode using electrospray ionisation (ESI) from methanol or acetonitrile. All reactions were performed under an atmosphere of dry, oxygen-free nitrogen in oven-dried glassware. Yields are given for isolated products showing one spot on a TLC plate and no impurities detectable in the NMR spectrum. The identity of the products prepared by different methods was checked by comparison of their NMR, IR, and MS data and by the TLC behavior. All chromatographic manipulations used silica gel as the adsorbent. Reactions were monitored by thin layer chromatography (TLC) on aluminum backed plates with Merck Kiesel 60 F254 silica gel. TLCs were either visualized by UV radiation at a wavelength of 254 nm, or stained by exposure to an ethanolic solution of phosphomolybdic acid or potassium permanganate aqueous solution, following by charring where appropriate. Purification by column chromatography was carried out using Merck Kiesel 60 H silica adsorbent. All solvents and reagents for the reactions were of reagent grade and were dried and distilled under nitrogen immediately before use as follows: tetrahydrofuran was distilled under a nitrogen atmosphere from the sodium/benzophenone ketyl radical, propionitrile and dichloromethane were freshly distilled from CaH₂. Petroleum ether refers to the fraction boiling in the range 40–60 °C. Synthesis of Z-crotyltrichlorosilane **10** and catalyst **12** was described by us earlier,^[5b] literature protocols were used for synthesising known aldehydes **11a**^[13] and **16**^[14].

Asymmetric synthesis of (–)-erogorgiaene (**1**) and its C-11 epimer (**1a**) by a revised strategy.

(E)-3-(2-bromo-3-methylphenyl)-2-methylacrylaldehyde (11c). A 2M solution of NaOH (1.9 mL, 3.8 mmol, 1 eq.) was added dropwise to a solution of benzaldehyde **16** (750 mg, 3.8 mmol, 1 eq.) and propanal (0.50 mL, 5.7 mmol, 1.5 eq.) in MeOH (15 mL), at rt. The reaction was allowed to stir at 50 °C overnight. After that time it was quenched with

NH₄Cl and then extracted with DCM (3 x 50 mL). The combined organic layers were dried over Na₂SO₄ and concentrated *in vacuo* to yield cinnamaldehyde **11b** (630 mg, 70% yield). ¹H NMR (400 MHz, CDCl₃): δ _H 1.92 (s, 3H); 2.47 (s, 3H); 7.20–7.22 (m, 1H); 7.25–7.27 (m, 2H); 7.50 (s, 1H); 9.69 (s, 1H); ¹³C NMR (100 MHz, CDCl₃): δ _C 10.7 (CH₃), 23.6 (CH₃), 126.6 (C), 126.7 (CH), 127.8 (CH), 131.2 (CH), 135.5 (C), 139.1 (C), 139.5 (C), 149.1 (CH), 195.3 (CH); IR: ν 3049, 2954, 2922, 2819, 2710, 1686, 1629, 1572, 1445, 14411, 1385, 1355, 1264, 1186, 1027, 1005, 935, 866, 834, 789, 771, 723, 681 cm⁻¹; HRMS (ESI) 239.0054 and 241.0034 (C₁₁H₁₁⁷⁹BrO and C₁₁H₁₁⁸¹BrO [M+H]⁺ requires 239.0066 and 241.0046).

(–)-(3R,4S,E)-1-(2-bromo-3-methylphenyl)-2,4-dimethylhexa-1,5-dien-3-ol (9c). A 100 mL round bottom flask fitted with a magnetic stirring bar was flame-dried, evacuated and backfilled with N₂. The flask was charged with 30 mL of dry propionitrile and successively with a solution of bis-*N*-oxide (–)-**12** (80.4 mg, 0.1 mmol, 2 mol%), aldehyde **11b** (1.25 g, 5.2 mmol, 1 eq.) and Hünig's base (1.8 mL, 10.3 mmol, 2 eq.) in 1 mL of dry propionitrile each. The solution was cooled at –60 °C and a solution of Z-crotyltrichlorosilane **10** (1.7 g, 8.9 mmol, 1.7 eq.) in 1 mL of dry propionitrile was added dropwise to the reaction mixture. The mixture was stirred at this temperature for two days. After that time, it was quenched with a saturated solution of NaHCO₃ (10 mL), washed with 20 mL of NaHCO₃ and extracted with DCM (3 x 60 mL). Combined organic layers were dried over Na₂SO₄ and solvent was removed *in vacuo*. Flash chromatography on silica with a gradient eluent system (100% hexane to 95:5, hexane/ethyl acetate) afforded the target homoallylic alcohol (–)-**9b** as a colourless oil (1.10 g, 72 % yield). ¹H NMR (400 MHz, CDCl₃): δ _H 1.13 (d, *J* = 6.8 Hz, 3H); 1.70 (s, 3H); 1.73 (s_{broad}, 1H); 2.43 (s, 3H); 2.53 (q, *J* = 6.8 Hz, 1H); 4.08 (d, *J* = 6.4 Hz, 1H); 5.06–5.15 (m, 2H); 5.83–5.91 (m, 1H); 6.53 (s, 1H); 7.04–7.06 (m, 1H); 7.11–7.17 (m, 2H); ¹³C NMR (100 MHz, CDCl₃): δ _C 13.9 (CH₃), 14.4 (CH₃), 23.7 (CH₃), 41.4 (CH), 80.0 (CH), 114.7 (CH₂), 126.3 (CH), 126.7 (C), 127.1 (CH), 128.2 (CH), 128.9 (CH), 138.3 (C), 138.3 (C), 139.5 (C), 140.9 (CH); IR: ν 3412, 3069, 2975, 2923, 2870, 1640, 1586, 1463, 1449, 1401, 1379, 1228, 1107, 1024, 913, 791, 767, 725 cm⁻¹; HRMS (ESI) 400.9666 and 402.9651 (C₁₅H₁₉⁷⁹BrO and C₁₅H₁₉⁸¹BrO [M+Ag]⁺ requires 400.9665 and 402.9644); Chiral HPLC (Chiralpak IA-3, hexane/2-propanol = 98:2, 0.75 mL/min, UV detection at 225 nm) showed 99:1 er (*t*_R = 19.3 min (major), *t*_S = 20.8 min (minor)). [α]_D²⁵ = –27 (c 6.5, CHCl₃).

In a racemic variant, Z-crotyltrichlorosilane **9c** (1.0 g, 5.00 mmol) was added to a solution of **11c** (75 mg, 0.31 mmol, 1 eq.) in DMF (5 mL) and Hünig's base (108 μ L, 0.62 mmol, 2 eq.) under N₂ atmosphere. The reaction mixture was stirred and monitored by TLC until completion. Then it was quenched with saturated NaHCO₃ (10 mL) and extracted with ethyl acetate (3 x 20 mL). The organic layers were dried over Na₂SO₄ and concentrated *in vacuo*. Flash chromatography on silica with a gradient eluent system (100% hexane to 95:5, hexane/ethyl acetate) afforded the target homoallylic alcohol (\pm)-**9c** as a colourless oil (78 mg, 85 %).

(2E,4R,5S,7E)-ethyl 5-(2-bromo-3-methylphenyl)-4-methylnona-2,7-dienoate (15c). A 30% suspension of KH in mineral oil (2.2 g, 55.5 mmol, 15 eq.) was washed with anhydrous DME (3 x 10 mL). Anhydrous DME (45 mL) was added to the KH. Alcohol (–)-**9c** (1.10 g, 3.7 mmol, 1 eq.) and 18-crown-6 (0.97 g, 3.7 mmol, 1 eq.) were successively added to the solution and the mixture was stirred for two days at rt. The solution was cooled to –78 °C and quenched with a 1M solution of 2,4,6-tri-*tert*-butylphenol in dry DME (7eq). The solution was poured into a saturated solution of NH₄Cl (50 mL) and extracted with DCM (3 x 50 mL). The combined organic layers were dried over Na₂SO₄ and concentrated *in vacuo*. The resulting crude aldehyde obtained as a yellow oil was used immediately in the next step.

LiCl (186.5 mg, 4.44 mmol, 1.2 eq.) was weighted in a 50 mL flame-dried round bottom flask. The flask was evacuated and backfilled with N₂. Dry CH₃CN (10 mL) was added followed by a solution of Hünig's base (0.8 mL, 4.44 mmol, 1.2 eq.) and triethyl phosphonoacetate (0.9 mL, 4.44 mmol, 1.2 eq.) in 2 mL of dry CH₃CN each. After stirring the mixture for 5 minutes, the crude aldehyde (3.7 mmol, 1 eq.) in 2 mL of anhydrous CH₃CN was added dropwise. The reaction mixture was stirred overnight at rt and after that time it was quenching with a saturated solution of NH₄Cl (20 mL). The mixture was extracted with DCM (3 x 50 mL), the organic phases dried over Na₂SO₄ and the solvents removed under reduced pressure. The crude mixture was purified by flash column chromatography on *silica* eluting with a gradient system (100% petroleum ether to 5:1 petroleum ether/ethyl acetate) to yield **15c** (dr 2:1) as a colourless oil (1.1 g, 81% yield over two steps).

Major isomer (taken as a mixture): ¹H NMR (400 MHz, CDCl₃): δ_H 1.01 (d, *J* = 6.8 Hz, 3H); 1.27 (t, *J* = 6.8 Hz, 3H); 1.53 (d, 6.4 Hz, 3H); 2.37-3.33 (m, 2H); 2.43 (s, 3H); 2.68-2.71 (m, 1H); 3.49-3.53 (m, 1H); 4.14-4.23 (m, 2H); 5.11-5.41 (m, 2H); 5.74 (d, *J* = 15.6 Hz, 1H); 6.92-7.01 (m, 2H); 7.06-7.17 (m, 2H).

Minor isomer (taken as a mixture): ¹H NMR (400 MHz, CDCl₃): δ_H 0.91 (d, *J* = 6.4 Hz, 3H); 1.31 (t, *J* = 7.2 Hz, 3H); 1.49 (d, *J* = 6.4 Hz, 3H); 2.37-3.33 (m, 2H); 2.44 (s, 3H); 2.49-2.53 (m, 1H); 3.38-3.41 (m, 1H); 4.14-4.23 (m, 2H); 5.11-5.41 (m, 2H); 5.82 (d, *J* = 14.8 Hz, 1H); 6.92-7.01 (m, 2H); 7.06-7.17 (m, 2H).

(*R,E*)-ethyl 4-((1*S,4R*)-8-bromo-4,7-dimethyl-1,2,3,4-tetrahydronaphthalen-1-yl)pent-2-enoate (16c). Methanesulfonic acid (0.37 mL, 5.7 mmol, 3 eq.) was added to a solution of **15c** (693 mg, 1.9 mmol, 1 eq.) in CHCl₃ (3 mL) and the reaction mixture was stirred at 40 °C for two days. The reaction was quenched with saturated NaHCO₃ (5 mL) and extracted with CH₂Cl₂ (3 x 20 mL). The combined organic layers were dried over Na₂SO₄ and concentrated *in vacuo*. Purification by column chromatography on *silica* (eluting with 95:5 petroleum ether/ethyl acetate) gave the *title compound 16c* (dr 2:1) as a pale yellow oil (570 mg, 83% yield);

Major isomer (taken as a mixture): ¹H NMR (400 MHz, CDCl₃): δ_H 0.89 (d, *J* = 6.8 Hz, 3H); 1.21-1.31 (m, 6H); 1.74-1.95 (m, 4H); 2.38 (s, 3H); 2.74-2.79 (m, 1H); 3.04-3.09 (m, 1H); 3.49-3.53 (m, 1H); 4.12-4.4.21 (m, 2H); 5.74 (dd, *J* = 16, 1.6 Hz, 1H); 7.06-7.12 (m, 3H); ¹³C NMR (100 MHz, CDCl₃): δ_C 14.2 (CH₃), 14.2 (CH₃), 16.9 (CH₃), 21.5 (CH₂), 22.3 (CH), 24.2 (CH₃), 29.2 (CH₂), 32.2 (CH), 39.6 (CH), 41.7 (CH), 60.1 (CH₂), 120.0 (CH), 125.4 (CH), 128.2 (CH), 136.1 (C), 137.8 (C), 143.4 (C), 153.3 (CH), 166.8 (C).

Minor isomer (taken as a mixture): ¹H NMR (400 MHz, CDCl₃): δ_H 0.90 (d, *J* = 6.8 Hz, 3H); 1.15 (d, *J* = 7.2 Hz, 3H); 1.21-1.31 (m, 3H); 1.74-1.95 (m, 4H); 2.41 (s, 3H); 2.74-2.79 (m, 1H); 2.97-3.00 (m, 1H); 3.40-3.44 (m, 1H); 4.12-4.21 (m, 2H); 5.64 (dd, *J* = 15.6, 1.6 Hz, 1H), 6.79 (dd, *J* = 15.6, 6.4 Hz, 1H); 7.06-7.07 (m, 2H); ¹³C NMR (100 MHz, CDCl₃): δ_C 14.2 (CH₃), 14.3 (CH₃), 16.9 (CH₃), 21.1 (CH₂), 22.6 (CH), 24.2 (CH₃), 28.5 (CH₂), 31.8 (CH), 40.0 (CH), 42.7 (CH), 60.0 (CH₂), 120.0 (CH), 125.7 (CH), 128.4 (CH), 136.0 (C), 138.1 (C), 143.1 (C), 152.6 (CH), 166.6 (C).

(-)-(R)-4-((1*S,4R*)-4,7-dimethyl-1,2,3,4-tetrahydronaphthalen-1-yl)pentan-1-ol (7) and **(-)-(S)-4-((1*S,4R*)-4,7-dimethyl-1,2,3,4-tetrahydronaphthalen-1-yl)pentan-1-ol (7a).** A solution of unsaturated ester **16c** (458 mg, 1.36 mmol, 1 eq.) in MeOH (35 mL) was circulated for 4 h through an H-Cube reactor (set at 2.0 mL/min, 20 bar, 40 °C) equipped with a cartridge containing catalyst 10% Pd/C. Analysis of an aliquote taken from the mixture showed that hydrogenation of the double

bond and debromination were complete. The solvent was removed *in vacuo*, the residue was dissolved in anhydrous CH₂Cl₂ (20 mL) and cooled to -78 °C. DIBAL, 1M in hexanes (2.2 mL, 2.2 mmol), was added and the solution was allowed to warm to room temperature and stirred overnight. The reaction was quenched with MeOH (10 mL), 5M HCl (15 mL) and extracted with CH₂Cl₂ (3x30 mL). The organic layer were combined, dried over Na₂SO₄ and concentrated *in vacuo*. The crude reaction mixture was filtered through a pad of silica. Removal of solvent *in vacuo* afforded product **7** (dr 2:1) as a colourless oil (235 mg, 70 %). The diastereoisomers **11-epi-7a** and **7** were separated in order of elution by preparative HPLC (Dynmax-60Å, hexane/2-propanol = 98:2, 5 mL/min, UV detection at 220 nm).

Major isomer (-)-7 (104 mg): ¹H NMR (400 MHz, CDCl₃): δ_H 0.65 (d, *J* = 5.2 Hz, 3H); 1.26 (d, *J* = 5.6 Hz, 3H); 1.31-1.38 (m, 3H); 1.61-1.70 (m, 3H); 1.78-1.84 (m, 1H); 1.90-1.95 (m, 1H); 2.13-2.16 (m, 1H); 2.30 (s, 3H); 2.71-2.75 (m, 1H); 2.87-2.91 (m, 1H); 3.70 (t, *J* = 5.2 Hz, 2H); 6.95 (d, *J* = 6.4 Hz, 1H); 7.03 (s, 1H); 7.14 (d, *J* = 6 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃): δ_C 14.5 (CH₃), 21.1 (CH₃), 21.5 (CH₂), 21.8 (CH₃), 31.1 (CH₂), 31.6 (CH₂), 32.8 (CH), 37.1 (CH), 41.6 (CH), 63.4 (CH₂), 126.0 (CH), 126.4 (CH), 128.0 (CH), 134.7 (C), 139.6 (C), 140.3 (C); IR: ν 3343, 2928, 2869, 1497, 1455, 1376, 1058, 812 cm⁻¹; HRMS (ESI) 353.1037 (C₁₇H₂₆O¹⁰⁷Ag [M]⁺ required 353.1029). [α]_D²⁵ = -54.5 (c 1.02, CHCl₃); lit. for (+)-**7** [α]_D²⁵ = +36.5 (c 0.81, CHCl₃).^[5c]

Minor isomer (-)-7a (65 mg): ¹H NMR (400 MHz, CDCl₃): δ_H 1.03 (d, *J* = 6.8 Hz, 3H); 1.28 (d, *J* = 6.8 Hz, 3H); 1.33-1.39 (m, 2H); 1.47 (s_{broad}, 1H); 1.55-1.60 (m, 3H); 1.89-1.94 (m, 2H); 2.08-2.13 (m, 1H); 2.32 (s, 3H); 2.75-2.81 (m, 2H); 3.48-3.56 (m, 2H); 6.96 (d, *J* = 8 Hz, 1H); 7.04 (s, 1H); 7.15 (d, *J* = 8 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃): δ_C 140.3 (C), 139.6 (C), 134.6 (C), 128.2 (CH), 126.8 (CH), 126.2 (CH), 63.1 (CH₂), 43.7 (CH), 36.9 (CH), 32.7 (CH), 31.2 (CH₂), 31.1 (CH₂), 27.3 (CH₂), 22.2 (CH₂), 22.1 (CH₃), 21.1 (CH₃), 18.2 (CH₃); IR: ν 3343, 2928, 2869, 1497, 1455, 1376, 1058, 812 cm⁻¹; HRMS (ESI) 353.1037 (C₁₇H₂₆O¹⁰⁷Ag [M]⁺ required 353.1029). [α]_D²⁵ = -32.9 (c 0.41, CHCl₃). [α]_D²⁵ = -54.5 (c 1.02, CHCl₃); lit. for (+)-**7** [α]_D²⁵ = +36.5 (c 0.81, CHCl₃).^[5c]

(-)-(1*R,4S*)-1,6-dimethyl-4-((*R*)-6-methylhept-5-en-2-yl)-1,2,3,4-tetrahydronaphthalene. (-)-Erogorgiaene (1)

Dess-Martin periodinane (1.5 eq.) was weighted in a 25 mL flame-dried round bottom flask. The vessel was evacuated and backfilled with N₂ before being charged with alcohol **7** (104 mg, 1 eq.) in 5 mL of DCM. The reaction was allowed to stir at room temperature for 2 hours. The solvent was partially removed *in vacuo* and the crude mixture was directly purified by flash column chromatography on *silica* using a 95:5 mixture of petroleum ether/ethyl acetate as an eluent. After removal of solvent *in vacuo*, the resulting aldehyde as pale yellow oil (86 mg, 84%) was used in the next step.

*n*BuLi, 1.6M solution in hexane (0.5 mL, 0.81 mmol, 3 eq.), was added dropwise to a solution of isopropyltriphenylphosphonium iodide (385.2 mg, 0.89 mmol) in anhydrous THF (3 mL) at 0 °C under N₂. The mixture was stirred for 1 h and then a solution of the aliphatic aldehyde from the step above (65 mg, 1.3 mmol) in THF (15 mL) was added. The solution was allowed to stir at the same temperature for 30 minutes before warming to rt and heating at reflux for a further 2 h. After cooling, the reaction mixture was quenched with saturated NH₄Cl (5 mL) and extracted with DCM (3 x 20 mL). The organic layers were combined, dried over Na₂SO₄ and concentrated *in vacuo*. Purification by column chromatography on *silica* (10:1 petroleum ether/ethyl acetate) gave (-)-erogorgiaene **1** as a colourless oil (58 mg, 80 %, or 67% over two steps). ¹H NMR (400 MHz, CDCl₃): δ_H 0.66 (d, *J* = 6.8 Hz, 3H); 1.29 (d, *J* = 6.8 Hz, 3H); 1.32-1.41 (m, 2H); 1.45-1.57 (m, 2H); 1.67 (s, 3H); 1.75 (s, 3H); 1.80-1.87 (m, 1H);

1.91-1.97 (m, 1H); 2.04-2.19 (m, 3H); 2.33 (s, 3H); 2.72-2.76 (m, 1H); 2.88-2.93 (m, 1H); 5.20 (tm, $J = 6.8$ Hz, 1H); 6.97 (d, $J = 8$ Hz, 1H); 7.05 (s, 1H); 7.16 (d, $J = 8$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3): δ_{C} 14.5 (CH_3), 17.7 (CH_3), 21.1 (CH_3), 21.5 (CH_2), 21.8 (CH_3), 25.7 (CH_3), 26.3 (CH_2), 31.8 (CH_2), 32.8 (CH), 35.2 (CH_2), 36.9 (CH), 41.4 (CH), 124.9 (CH), 126.0 (CH), 126.4 (CH), 128.1 (CH), 131.2 (C), 134.7 (C), 139.9 (C), 140.4 (C); IR: ν 2959, 2924, 2853, 1613, 1497, 1452, 1376, 1320, 1109, 1039, 983, 881, 811 cm^{-1} ; HRMS (ESI) 377.1390 and 379.1386 ($\text{C}_{20}\text{H}_{30}^{107}\text{Ag}$ [M+Ag] $^+$ requires 377.1393 and $\text{C}_{20}\text{H}_{30}^{109}\text{Ag}$ [M+Ag] $^+$ requires 379.1390). $[\alpha]_{\text{D}}^{25} = -25.5$ (c 0.52, CHCl_3); lit. $[\alpha]_{\text{D}}^{25} = -28.0$ (c 0.05, CHCl_3);^[2] lit. for (+)-**1**: $[\alpha]_{\text{D}}^{25} = +24.4$ (c 3.20, CHCl_3);^[1a] $[\alpha]_{\text{D}}^{25} = +23.2$ (c 0.75, CHCl_3);^[3c] $[\alpha]_{\text{D}}^{25} = +69.1$ (c 0.77, DCM);^[4b] $[\alpha]_{\text{D}}^{25} = +40.6$ (c 0.14, CHCl_3);^[5a] $[\alpha]_{\text{D}}^{25} = +21.4$ (c 0.14, CHCl_3).^[5c]

(-)-(1R,4S)-1,6-dimethyl-4-((S)-6-methylhept-5-en-2-yl)-1,2,3,4-tetrahydronaphthalene (1a).

Diastereoisomer (-)-**1a** was obtained from (-)-**7a** using the same sequence as described for (-)-**1** as a colourless oil (15 mg, 30% from **7a**): ^1H NMR (400 MHz, CDCl_3): δ_{H} 1.00 (d, $J = 6.8$ Hz, 3H); 1.23-1.27 (m, 7H); 1.55 (s, 3H); 1.66 (s, 3H); 1.80-2.00 (m, 4H); 2.07-2.12 (m, 2H); 2.30 (s, 3H); 2.73-2.77 (m, 2H); 4.98 (t, $J = 7.4$ Hz, 1H); 6.94 (m, 1H); 7.03 (s, 1H); 7.13 (d, $J = 8$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3): δ_{C} 17.6 (CH_3), 18.1 (CH_3), 21.1 (CH_3), 22.2 (CH_3), 22.4 (CH_2), 25.7 (CH_3), 26.3 (CH_2), 31.4 (CH_2), 31.5 (CH_2), 32.7 (CH), 36.2 (CH), 43.8 (CH), 125.0 (CH), 126.1 (CH), 126.7 (CH), 128.4 (CH), 131.2 (C), 134.5 (C), 139.7 (C), 140.2 (C). IR: ν 2959, 2924, 2853, 1613, 1497, 1452, 1376, 1320, 1109, 1039, 983, 881, 811 cm^{-1} ; HRMS (ESI) 377.1390 and 379.1386 ($\text{C}_{20}\text{H}_{30}^{107}\text{Ag}$ [M+Ag] $^+$ requires 377.1393 and $\text{C}_{20}\text{H}_{30}^{109}\text{Ag}$ [M+Ag] $^+$ requires 379.1390). $[\alpha]_{\text{D}}^{25} = -25.3$ (c = 0.3, CHCl_3); lit. for (+)-**1a**: $[\alpha]_{\text{D}}^{22} = +55.2$ (c 0.83, DCM).^[4b]

Acknowledgements

The authors thank Russian Science Foundation for grant 15-13-00092 and the Loughborough University for studentships to CAIP and PSOH.


Keywords: allylation • asymmetric catalysis • rearrangement • stereoselectivity • total synthesis

- [1] a) A. D. Rodriguez, C. Ramirez, *J. Nat. Prod.* **2001**, *64*, 100-102; b) M. Donia, M. T. Hamann, *Lancet. Infect. Dis.* **2003**, *3*, 338-348; c) C. P. Ndi, S. J. Semple, H. J. Griesser, S. M. Pyke, M. D. Barton, *Phytochemistry* **2007**, *68*, 2684-2690; d) H. Correa, F. Aristizabal, C. Duque, R. Kerr, *Marine drugs* **2011**, *9*, 334-343; e) O. F. Anakok, C. P. Ndi, M. D. Barton, H. J. Griesser, S. J. Semple, *J. Appl. Microbiol.* **2012**, *112*, 197-204; f) E. C. Barnes, A. M. Kavanagh, S. Ramu, M. A. Blaskovich, M. A. Cooper, R. A. Davis, *Phytochemistry* **2013**, *93*, 162-169; g) E. Sansinenea, A. Ortiz, *Curr. Org. Synth.* **2014**, *11*, 545-591.
- [2] S. A. Kolesnikova, A. I. Kalinovsky, S. N. Fedorov, L. K. Shubina, V. A. Stonik, *Phytochemistry* **2006**, *67*, 2115-2119.
- [3] a) J. M. H. Lu, M. V. Perkins, H. J. Griesser, *Tetrahedron* **2013**, *69*, 6468-6473; b) J. S. Yadav, A. K. Basak, P. Srihari, *Tetrahedron Lett.*

- 2007**, *48*, 2841-2843; c) J. S. Yadav, B. Thirupathaiah, A. Al Khazim Al Ghamdi, *Eur. J. Org. Chem.* **2012**, 2072-2076.
- [4] a) F. Dehmel, J. Lex, H. G. Schmalz, *Org. Lett.* **2002**, *4*, 3915-3918; b) T. G. Eilford, S. Nave, R. P. Sonawane, V. K. Aggarwal, *J. Am. Chem. Soc.* **2011**, *133*, 16798-16801; c) M. Harmata, X. Hong, *Tetrahedron Lett.* **2005**, *46*, 3847-3849; d) M. Harmata, X. Hong, P. R. Schreiner, *J. Org. Chem.* **2008**, *73*, 1290-1296; e) W. Lolsberg, S. Werle, J. M. Neudorff, H. G. Schmalz, *Org. Lett.* **2012**, *14*, 5996-5999.
- [5] a) R. R. Cesati lii, J. De Armas, A. H. Hoveyda, *J. Am. Chem. Soc.* **2004**, *126*, 96-101; b) H. M. L. Davies, X. Dai, *Tetrahedron* **2006**, *62*, 10477-10484; c) H. M. L. Davies, A. M. Walji, *Angew. Chem.* **2005**, *117*, 1761-1763; *Angew. Chem. Int. Ed.* **2005**, *44*, 1733-1735; d) P. S. O'Hara, C. A. Incerti-Pradillos, M. A. Kabeshov, S. A. Shipilovskikh, A. E. Rubtsov, M. R. J. Elsegood, A. V. Malkov, *Chem. Eur. J.* **2015**, *21*, 4551-4555; e) X. Yu, F. Su, C. Liu, H. Yuan, S. Zhao, Z. Zhou, T. Quan, T. Luo, *J. Am. Chem. Soc.* **2016**, *138*, 6261-6270.
- [6] a) S. E. Denmark, J. Fu, M. J. Lawler, *J. Org. Chem.* **2006**, *71*, 1523-1536; b) A. Kadlčíková, I. Valterová, L. Ducháčková, J. Roithová, M. Kotora, *Chem. Eur. J.* **2010**, *16*, 9442-9445; c) A. V. Malkov, M. Barlóg, Y. Jewkes, J. Mikušek, P. Kočovský, *J. Org. Chem.* **2011**, *76*, 4800-4804; d) P. Koukal, J. Ulč, D. Nečas, M. Kotora, *Eur. J. Org. Chem.* **2016**, 2110-2114.
- [7] a) L. A. Paquette, *Angew. Chem.* **1990**, *102*, 642-660; *Angew. Chem. Int. Ed.* **1990**, *29*, 609-626; b) L. A. Paquette, G. D. Maynard, *Angew. Chem.* **1991**, *103*, 1392-1394; *Angew. Chem. Int. Ed.* **1991**, *30*, 1368-1370; c) L. A. Paquette, G. D. Maynard, *J. Am. Chem. Soc.* **1992**, *114*, 5018-5027; d) A. P. Rutherford, R. C. Hartley, *Tetrahedron Lett.* **2000**, *41*, 737-741; e) H. T. Mamdani, R. C. Hartley, *Tetrahedron Lett.* **2000**, *41*, 747-749; f) K. Tomooka, S.-Y. Wei, T. Nakai, *Chem. Lett.* **1991**, 43-46. For recent overviews, see g) E. A. Ildardi, C. E. Stivala, A. Zakarian, *Chem. Soc. Rev.* **2009**, *38*, 3133-3148; h) H. M. Davies, Y. Lian, *Acc. Chem. Res.* **2012**, *45*, 923-935.
- [8] R. Appelbe, M. Casey, A. Dunne, E. Pascarella, *Tetrahedron Lett.* **2003**, *44*, 7641-7644.
- [9] For details of conformational search for the enol form of **8c**, see ESI.
- [10] a) L. Czollner, W. Frantsits, B. Küenburg, U. Hedenig, J. Fröhlich, U. Jordis, *Tetrahedron Lett.* **1998**, *39*, 2087-2088; b) B. Küenburg, L. Czollner, J. Fröhlich, U. Jordis, *Org. Process Res. Dev.* **1999**, *3*, 425-431.
- [11] M. A. Blanchette, W. Choy, J. T. Davis, A. P. Essensfeld, S. Masamune, W. R. Roush, T. Sakai, *Tetrahedron Lett.* **1984**, *25*, 2183-2186.
- [12] a) R. Escarcena, J. Perez-Meseguer, E. del Olmo, B. Alanis-Garza, E. Garza-Gonzalez, R. Salazar-Aranda, N. W. de Torres, *Molecules* **2015**, *20*, 7245-7262; b) G. M. Molina-Salinas, V. M. Rivas-Galindo, S. Said-Fernandez, D. C. Lankin, M. A. Munoz, P. Joseph-Nathan, G. F. Pauli, N. Waksman, *J. Nat. Prod.* **2011**, *74*, 1842-1850; c) J. Perez-Meseguer, E. del Olmo, B. Alanis-Garza, R. Escarcena, E. Garza-Gonzalez, R. Salazar-Aranda, A. San Feliciano, N. W. de Torres, *Bioorg. Med. Chem.* **2012**, *20*, 4155-4163.
- [13] X. Fan, H. Lv, Y. H. Guan, H. B. Zhu, X. M. Cui, K. Guo, *Chem. Commun.* **2014**, 50, 4119-4122.
- [14] J. Carreras, G. Gopakumar, L. G. Gu, A. Gimeno, P. Linowski, J. Petuskova, W. Thiel, M. Alcarazo, *J. Am. Chem. Soc.* **2013**, *135*, 18815-18823.

Entry for the Table of Contents (Please choose one layout)

FULL PAPER



(-)-Erogorgiaene

C11-epimer

A short, 9-step, highly enantioselective synthesis of (-)-erogorgiaene and its C11 epimer is reported. The key stereochemistry control-

ling steps involve catalytic asymmetric crotylation, anionic oxy-Cope rearrangement and cationic cyclisation. (-)-Erogorgiaene exhibited promising antitubercular activity against multidrug resistant strains of *M. tuberculosis*

Diterpenes*

Celia A. Incerti-Pradillos, Mikhail A. Kabeshov, Paul S. O'Hora, Sergei A. Shipilovskikh, Aleksandr E. Rubtsov, Vera A. Drobkova, Svetlana Yu. Balandina, and Andrei V. Malkov*

Page No. – Page No.

Asymmetric Total Synthesis of (-)-Erogorgiaene and Its C11 Epimer and Investigation of Their Antimycobacterial Activity