

Loughborough University Institutional Repository

A single day of excessive dietary fat intake reduces whole-body insulin sensitivity: the metabolic consequence of binge eating

This item was submitted to Loughborough University's Institutional Repository by the/an author.

Citation: PARRY, S.A. ... et al, 2017. A single day of excessive dietary fat intake reduces whole-body insulin sensitivity: the metabolic consequence of binge eating. *Nutrients*, 9 (8), 818.

Additional Information:

- This is an Open Access Article. It is published by MDPI under the Creative Commons Attribution 4.0 International Licence (CC BY). Full details of this licence are available at: <http://creativecommons.org/licenses/by/4.0/>

Metadata Record: <https://dspace.lboro.ac.uk/2134/26116>

Version: Published

Publisher: MDPI AG © The Authors

Rights: This work is made available according to the conditions of the Creative Commons Attribution 4.0 International (CC BY 4.0) licence. Full details of this licence are available at: <http://creativecommons.org/licenses/by/4.0/>

Please cite the published version.

Article

A Single Day of Excessive Dietary Fat Intake Reduces Whole-Body Insulin Sensitivity: The Metabolic Consequence of Binge Eating

Siôn A. Parry¹, Rachel M. Woods¹, Leanne Hodson²  and Carl J. Hulston^{1,*}

¹ School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, Leicestershire LE11 3TU, UK; S.Parry@lboro.ac.uk (S.A.P.); R.M.Woods@lboro.ac.uk (R.M.W.)

² Oxford Centre for Diabetes, Endocrinology and Metabolism, University of Oxford, Churchill Hospital, Oxford OX3 7LE, UK; leanne.hodson@ocdem.ox.ac.uk

* Correspondence: c.j.hulston@lboro.ac.uk; Tel.: +44-150-922-6449

Received: 5 June 2017; Accepted: 26 July 2017; Published: 29 July 2017

Abstract: Consuming excessive amounts of energy as dietary fat for several days or weeks can impair glycemic control and reduce insulin sensitivity in healthy adults. However, individuals who demonstrate binge eating behavior overconsume for much shorter periods of time; the metabolic consequences of such behavior remain unknown. The aim of this study was to determine the effect of a single day of high-fat overfeeding on whole-body insulin sensitivity. Fifteen young, healthy adults underwent an oral glucose tolerance test before and after consuming a high-fat (68% of total energy), high-energy (78% greater than daily requirements) diet for one day. Fasting and postprandial plasma concentrations of glucose, insulin, non-esterified fatty acids, and triglyceride were measured and the Matsuda insulin sensitivity index was calculated. One day of high-fat overfeeding increased postprandial glucose area under the curve (AUC) by 17.1% ($p < 0.0001$) and insulin AUC by 16.4% ($p = 0.007$). Whole-body insulin sensitivity decreased by 28% ($p = 0.001$). In conclusion, a single day of high-fat, overfeeding impaired whole-body insulin sensitivity in young, healthy adults. This highlights the rapidity with which excessive consumption of calories through high-fat food can impair glucose metabolism, and suggests that acute binge eating may have immediate metabolic health consequences for the individual.

Keywords: glucose; insulin; high-fat; overfeeding; glycemic control; type 2 diabetes mellitus

1. Introduction

Binge eating, which is defined as discreet periods of excessive food consumption that is not driven by hunger or metabolic need [1,2], is strongly associated with overweight and obesity [3–5], which are risk factors for developing insulin resistance and the metabolic syndrome [6]. Those who exhibit binge eating behavior are known to eat until they are uncomfortably full and then may or may not compensate for this increased energy intake, with the latter leading to a positive energy balance [1]. The etiology of binge eating and binge eating disorders (BED: the clinical manifestation of binge eating behavior [7]) has been extensively studied from a psychological perspective, with findings indicating that a number of stressors (e.g., interpersonal, ego-threatening, and work-related) are associated with increased food intake and inter-meal snacking [8,9]. However, the metabolic response to such behavior has received relatively little attention and, therefore, little is known regarding the metabolic consequences.

Binge eating has been associated with a selective increase in the intake of palatable foods (i.e., foods which are high in fat and/or sugar) [7]. This is important, as dietary composition appears to be a key mediating feature in the pathogenesis of metabolic disease. For example, high-fat diets

or the consumption of saturated fatty acid (SFA)-enriched diets have been associated with insulin resistance [10,11]. Results from the Kuopio, Aarhus, Naples, Wollongong and Uppsala (KANWU) study demonstrated that consumption of an SFA-rich diet for three months reduced insulin sensitivity in healthy adults when compared to a diet enriched in monounsaturated fatty acids (MUFA) [12]. Furthermore, when stratified by habitual total fat intake (above and below median), individuals with a higher fat intake (>37% total energy (TE)) exhibited reduced insulin sensitivity compared to those with a low intake (<37% TE) [13]. The mechanism by which a high total fat or SFA-diet causes insulin resistance remains unclear. One hypothesis is that adipose tissue dysfunction leads to an overabundance of circulating non-esterified fatty acid (NEFA) and insulin resistance ensues due to the accumulation of NEFA in non-adipose tissue organs, such as skeletal muscle and the liver [14]. Although work from the 1960s [15,16] supports the hypothesis as elevated NEFA concentrations were observed in obesity; recent reports are not in agreement [17]. Experimentally elevating plasma NEFA concentrations, via intravenous (iv) infusion of a lipid-heparin emulsion (Intralipid), rapidly (within 6–8 h) reduces insulin-mediated glucose disposal in healthy, lean individuals [18–20]. This is associated with a sequential pattern of events starting with an increase in circulating triglyceride (TG) levels (<0.5 h), followed by a supraphysiological increase in circulating NEFA levels (~1–1.5 h), a rise in intramyocellular lipid (IMCL) content (~2.5 h), and finally a reduction in insulin sensitivity; this supports the concept that elevated NEFA levels are a key player in the development of insulin resistance. However, during iv lipid infusions, plasma NEFA concentrations are typically elevated to around 1500 $\mu\text{mol/L}$ or higher [17], which is in excess of the fasting concentrations reported for obese, insulin-resistant individuals (~400 $\mu\text{mol/L}$) [21], and individuals with poorly managed type 2 diabetes mellitus (T2DM) (~800 $\mu\text{mol/L}$) [22].

Results from short-term (4–14 days), high-fat overfeeding interventions report impaired glycemic control [23–25] and decreased hepatic [26] and whole-body insulin sensitivity [27,28] despite unchanged or reduced plasma NEFA concentrations. Although these studies demonstrate that impairments in glycemic control occur after several days or weeks of consistent overconsumption, this model may not reflect the dietary practices of those who binge eat and consume a severe energy excess within a matter of hours. Although it remains unclear if episodes of binge eating have a negative impact on glucose metabolism, there is some evidence, albeit limited, to suggest that diet-induced impairments may occur very rapidly. Nowotny et al. [29] reported that oral administration of a single dose of soybean oil (100 mL), which is enriched with polyunsaturated fat (61% polyunsaturated, 23% monounsaturated, and 16% saturated), reduced whole-body insulin sensitivity (assessed by hyperinsulinemic-euglycemic clamp) to a comparable extent and within a similar time-frame (6 h post ingestion/infusion) as an energy and composition-matched iv lipid-heparin infusion. This occurred independent of plasma NEFA levels, which were elevated during the iv fat infusion but were unchanged following oral fat ingestion [29]. Insulin sensitivity was assessed 6–8 h after fat ingestion/infusion [29], and it is possible that the observed reduction in insulin sensitivity was a transient response related to the ongoing metabolism of fat; it would be of interest to determine if changes persist into the post-absorptive state and occur after the consumption of a diet more reflective of Western style eating patterns (i.e., SFA rather than polyunsaturated fatty acid (PUFA)-enriched). Therefore, in order to replicate excessive binge eating behavior, we undertook a pilot study in which whole-body insulin sensitivity was assessed in young, healthy, non-obese individuals after a single day of high-energy, high-fat, SFA-rich food intake.

2. Materials and Methods

2.1. Subjects

Fifteen healthy individuals were recruited for this study. All subjects were physically active (exercising at least three times per week for more than 30 min at a time), non-smokers, free from cardiovascular and metabolic disease, not taking any medication, and weight stable for at least

six months. The study was conducted according to the Declaration of Helsinki and was approved by the Loughborough University Ethical subcommittee for human participants (ethical approval number R13-P171). All subjects gave written informed consent.

2.2. Pre-Testing

Prior to the start of the study, subjects attended the laboratory for an initial assessment of baseline anthropometric characteristics (height, weight, and body mass index (BMI)) which were used to estimate resting energy expenditure (REE) using the calculations described by Mifflin et al. [30]. A standard correction for physical activity level (1.6 and 1.7 times REE for females and males, respectively) was applied in order to estimate total daily energy requirements. This information was then used to determine individual energy intakes for the one-day overfeeding period.

2.3. Experimental Design

After the pre-testing visit, subjects attended the laboratory for an oral glucose tolerance test (OGTT) and then continued their habitual food intake for six days. On the seventh day, subjects consumed the experimental diet that was provided to them. The experimental diet was designed to be high in fat (68% total energy) and provide an energy excess (+78% kJ). High-fat foods were specifically chosen because individuals tend to overconsume more readily with high-fat foods due to the greater palatability and the higher energy density of this macronutrient, and because individuals who binge eat are known to selectively increase their intake of such foods [7]. Individual diet plans were designed using NetWISP nutrition software (Tinuviel Software Ltd., Llanfechell, Anglesey, UK). All foods were purchased and prepared by the research team. Subjects were instructed to consume all food provided, and to avoid consuming additional food or nutritive beverages. Food intake followed a normal daily feeding pattern (i.e., breakfast, lunch, dinner, and snacks) and water intake was allowed ad libitum throughout the dietary intervention. An example diet plan for one subject can be viewed in the supplementary material online. No subjects reported any issues with dietary adherence. The next day (Day 8), subjects returned to the laboratory for a second OGTT.

2.4. Experimental Protocol

On the experimental days (before (Day 0) and after overfeeding (Day 8)), subjects reported to the laboratory between 07.00 and 09.00 h after an overnight fast of at least 10 h and having refrained from physical activity for 48 h. After being weighed, a 20-gauge Teflon catheter (Venflon, Becton, Dickinson, Plymouth, UK) was inserted into an antecubital vein of an arm to allow for repeated blood sampling during the 2 h OGTT. A baseline (fasted (Time 0)) blood sample was taken and then subjects consumed a 25% glucose solution (75 g of glucose dissolved in 300 mL of water). Blood samples were then taken at 15, 30, 45, 60, 90, and 120 min after glucose ingestion.

2.5. Blood Sampling

Whole blood was collected into pre-chilled, ethylenediaminetetraacetic acid (EDTA; 1.75 mg/mL)-treated tubes (Sarstedt, Leicester, UK) and immediately spun at 1750 g in a refrigerated centrifuge (4 °C) for 10 min to obtain plasma, which was then stored at −20 °C until analysis. For the collection of serum, whole-blood was collected into tubes containing a clotting catalyst (Sarstedt, Leicester, UK) and left at room temperature until complete clotting had occurred. Samples were then centrifuged at 1750 g in a refrigerated centrifuge (4 °C) for 10 min and serum collected and stored at −20 °C until analysis.

2.6. Analytical Procedures

Plasma samples were analyzed using commercially available spectrophotometric assays for glucose, TG (Glucose PAP CP A11A01668, Triglycerides CP A11A01640; Horiba Medical, Northampton,

UK), and NEFA (FA115; Randox, County Antrim, UK) concentrations using a semi-automatic analyzer (Pentra 400; Horiba Medical, Northampton, UK). Serum insulin concentrations were determined using an enzyme-linked immuno-sorbent assay (ELISA: EIA-2935, DRG instruments GmbH, Marburg, Germany).

2.7. Calculations

Plasma glucose and serum insulin concentrations obtained before and during the OGTT were used to determine whole-body insulin sensitivity using the Matsuda insulin sensitivity index (ISI):

$$ISI = \frac{10,000}{\sqrt{(FPG \times FSI) \times (\text{mean OGTT insulin concentration}) \times (\text{mean OGTT glucose concentration})}}$$

where FPG is the fasting plasma glucose concentration; FSI is the fasting serum insulin concentration; and 10,000 represents a constant that allows numbers ranging between 1 and 12 to be obtained. The square root conversion is used to correct the nonlinear distribution of values [31]. Area under the curve (AUC) for glucose and insulin was calculated using the trapezoidal rule.

2.8. Statistics

Data are presented as means \pm standard error of the mean (SEM). Statistical analysis was performed using SPSS (V21.0) for windows (SPSS Inc., Chicago, IL, USA). Fasting concentrations of glucose, insulin, NEFA, and TG before and after high-fat overfeeding were compared using a paired *t*-test, whereas the dynamic hormonal and metabolic responses to the OGTT were compared using a two-way (trial \times time) repeated measures analysis of variance (ANOVA) and Bonferroni post hoc analysis where appropriate. Statistical significance was set at $p < 0.05$.

3. Results

3.1. Diet Intervention

The estimated energy requirement, actual energy intake, macronutrient intake, and fatty acid composition of the one-day high-fat overfeeding intervention are shown in Table 1. An example diet plan for one of the subjects can be seen in the supplementary material available online.

Table 1. Energy and nutrient intakes.

	Estimated Energy Requirement	Experimental Energy Intake
Energy (kJ)	14,028 \pm 433	24,949 \pm 797 *
Carbohydrate (g)		192 \pm 6
Protein (g)		278 \pm 8
Fat (g)		449 \pm 15
Fatty acid composition (%)		
SFA		42 \pm 0.6
MUFA		40 \pm 0.4
PUFA		10 \pm 0.2

Data presented are means \pm SEM. $n = 15$. * $p < 0.05$ significantly different from estimated energy requirements. Abbreviations: SFA, total saturated fatty acids; MUFA, total monounsaturated fatty acids; PUFA, total polyunsaturated fatty acids.

3.2. Subject Characteristics

Fifteen insulin-sensitive (5.0 ± 0.5 Matsuda ISI) individuals (13 males and 2 females), with a mean BMI of 26.4 ± 1.1 kg/m² were recruited (Table 2). Subjects gained on average 0.85 ± 0.20 kg body mass following one-day of high-fat overfeeding ($p = 0.001$), resulting in a BMI increase of

$0.26 \pm 0.06 \text{ kg/m}^2$ ($p < 0.0001$) (Table 2). Whole-body insulin sensitivity (assessed by the Matsuda ISI) was reduced by 28% ($p = 0.001$) after high-fat overfeeding (Table 2). There was no change in fasting serum insulin or plasma TG concentrations following high-fat overfeeding (Table 2). Fasting plasma glucose concentrations tended to increase after high-fat overfeeding, although this did not reach significance ($p = 0.058$), whilst fasting plasma NEFA were significantly decreased ($p = 0.009$) (Table 2).

Table 2. Subject characteristics before and after one day of high-fat overfeeding.

	Before Overfeeding	After Overfeeding
Males/Females	13/2	
Age (years)	22.1 \pm 0.5	
Weight (kg)	86.0 \pm 3.2	86.8 \pm 3.2 *
BMI (kg/m ²)	26.4 \pm 1.1	26.6 \pm 1.1 *
Fasting plasma biochemical parameters and Matsuda ISI		
Glucose (mmol/L)	5.5 \pm 0.1	5.7 \pm 0.1
Insulin (pmol/L)	80 \pm 9	98 \pm 20
NEFA (mmol/L)	0.43 \pm 0.06	0.24 \pm 0.02 *
TG (mmol/L)	0.91 \pm 0.08	0.78 \pm 0.10
Matsuda ISI	5.0 \pm 0.5	3.6 \pm 0.4 *

Data presented are means \pm SEM. $n = 15$. * $p < 0.05$ significantly different from before overfeeding. Abbreviations: BMI, body mass index; TG, triglyceride; NEFA, non-esterified fatty acids; ISI, insulin sensitivity index.

3.3. Oral Glucose Tolerance Test

Plasma glucose and serum insulin concentrations increased in response to the OGTT, peaking 30–45 min after ingestion (Figure 1A,B). There was a significant trial \times time interaction ($p = 0.002$) for plasma glucose (Figure 1A) but not for serum insulin. Postprandial plasma glucose concentrations increased by 17.1% after overfeeding (AUC, from $785 \pm 35 \text{ mmol/L per 120 min}$ before overfeeding to $920 \pm 33 \text{ mmol/L per 120 min}$ after overfeeding, $p < 0.001$). Postprandial serum insulin concentrations increased by 16.4% after overfeeding (AUC, from $39,462 \pm 2840 \text{ pmol/L per 120 min}$ before overfeeding to $45,947 \pm 3396 \text{ pmol/L per 120 min}$ after overfeeding, $p = 0.007$). A significant ($p < 0.0001$) time \times trial interaction was noted for plasma NEFA (Figure 1C) despite plasma the NEFA AUC not differing between the two study days ($22 \pm 2 \text{ mmol/L per 120 min}$ before overfeeding and $24 \pm 2 \text{ mmol/L per 120 min}$ after overfeeding, $p = 0.468$). A significant ($p < 0.0001$) time \times trial interaction was also observed for plasma TG concentrations (Figure 1D), although average TG concentrations over the course of the postprandial period were not significantly different between study days ($0.85 \pm 0.06 \text{ mmol/L}$ before overfeeding and $0.81 \pm 0.07 \text{ mmol/L}$ after overfeeding, $p = 0.614$).




Figure 1. Cont.




Figure 1. Systemic plasma glucose (A), serum insulin (B), plasma non-esterified fatty acid (NEFA) (C), and plasma triglyceride (TG) (D) concentrations during a 2-h oral glucose tolerance test (OGTT) conducted before (pre) and after (post) one day of high-fat overfeeding (HFD). Data presented are means \pm standard error of the mean (SEM). $n = 15$. * $p < 0.05$ significant difference between trials at the annotated time point. # $p < 0.05$ denotes significant main effect of trial/high-fat overfeeding.

4. Discussion

Short-term (4–14 days) adherence to a high-fat, high-energy diet has previously been reported to impair glycemic control and reduce insulin sensitivity in healthy individuals [23–25,27,28]. The findings we report here from our pilot study build upon this work and suggest that a single day of high-fat, high-energy food consumption impairs whole-body insulin sensitivity; evidenced by a significant (28%) reduction in whole-body insulin sensitivity as calculated by Matsuda ISI. Although our dietary model was quite extreme, with energy intake being approximately 78% greater than estimated daily requirements, individuals who demonstrate binge eating behaviors and those with BED frequently consume abnormally large amounts of food over a short period of time (i.e., within a matter of hours) [1–3]. Our results suggest that even brief periods of excessive consumption of foods that are typical of a Western diet may lead to metabolic dysfunction.

In the present study, participants consumed an excess energy intake as well as a high proportion of SFA. We chose this dietary intervention as the availability of palatable foods (e.g., those higher in fat) has been reported to be a precursory stimuli for binge eating [32,33] and individuals are more likely to overconsume with fat-rich foods due to the higher energy density of this macronutrient. Due to the composition of the experimental diet, we are unable to determine whether it is excess energy, excess dietary fat, or a combination of both that negatively impact glucose metabolism. Whether feeding excess energy in the form of carbohydrates (particularly added sugars, which are also highly palatable) for a single day has the same effect remains unclear. However, overfeeding a carbohydrate-rich diet (40% increase in energy intake; 60% of energy from carbohydrate) for five days was found to elicit changes in skeletal muscle cellular signaling that are typically associated with increased insulin sensitivity (i.e., increased tyrosine phosphorylation of insulin receptor-1 (IRS-1) and increased phosphatidylinositol 3 (PI 3)-kinase activity) whilst high-fat overfeeding was associated with reductions in markers of skeletal muscle insulin sensitivity (i.e., increased serine phosphorylation of IRS-1 and increased total expression of p85 α) [34]. These data suggest that excessive consumption of dietary fat reduces whole-body insulin sensitivity, rather than a positive energy balance alone.

Our finding that one day of high-fat overfeeding reduces whole-body insulin sensitivity was associated with a significant (17.1%) increase in postprandial glucose AUC. Postprandial glucose homeostasis is regulated by a number of factors, including the appearance of ingested glucose, endogenous glucose production, and splanchnic and peripheral glucose uptake [35,36]. Studies in type 2 diabetic subjects demonstrate that impairments in each of these regulatory factors contribute to postprandial hyperglycemia [37–44]. As the present study was a pilot study, we measured insulin sensitivity at the whole-body level using an OGTT in combination with a validated insulin sensitivity

index (the Matsuda ISI [31]). Whilst this method is easy to perform, it does not allow us to determine the contributions of each of the factors that may contribute to elevated postprandial glucose levels. Furthermore, we did not measure additional factors that may influence data, including the neural and incretin hormone response to nutrient ingestion [45,46]. It would be of interest to perform a high-fat and/or high-carbohydrate overfeeding study where whole-body and tissue-specific insulin sensitivity was assessed using clamp techniques and stable-isotope tracers [47–49]. Based on the observations of Brons et al. [26], it is plausible that increased endogenous glucose production may underpin early diet-induced impairments in whole-body insulin sensitivity, although others suggest skeletal muscle (i.e., reduced glucose uptake) as the primary site of metabolic dysfunction [29,34,50].

In line with the discussion point above, the mechanisms underpinning the observed reduction in insulin sensitivity are yet to be elucidated. Elevated plasma NEFA concentrations have previously been implicated in the development of insulin resistance. In the present study, although fasting NEFA concentrations were reduced following high-fat overfeeding, and NEFA AUC was not significantly different between the two study days (which is in line with our previous observations following seven days of high-fat overfeeding [24]), we observed a reduction in markers of whole-body insulin sensitivity. When NEFA concentrations are elevated experimentally via Intralipid infusion, insulin sensitivity is rapidly (within 6–8 h) reduced in healthy individuals [18–20]. Moreover, it has been demonstrated that the ingestion of a single oral fat bolus reduced whole-body insulin sensitivity to a comparable degree, and within a corresponding time-frame (6 h post ingestion/infusion), as that seen with an energy- and composition-matched Intralipid infusion, despite divergent plasma NEFA responses [29]. The observed reduction in insulin sensitivity following both Intralipid infusion and oral fat ingestion appeared to be mediated by the increased activation of protein kinase C theta (PKC θ), which is suggested to impair insulin signaling and reduce insulin-mediated glucose uptake by inhibiting the normal tyrosine kinase cascade via phosphorylation of the counter-regulatory serine residue of insulin receptor substrate-1 (IRS-1) [51]. The reason for this apparent divergence between circulating NEFA concentrations and insulin resistance remains unclear. Whilst our findings are not directly comparable, they highlight that the relationship between circulating NEFA levels and insulin resistance is not yet fully understood.

Increased plasma TG concentrations are a characteristic feature of T2DM and the metabolic syndrome [52]. Pramfalk et al. [53] recently demonstrated that hyperinsulinemic individuals exhibit increased hepatic de novo lipogenesis and hypertriglyceridemia compared to those who were normoinsulinemic [53], which has been hypothesized to be due to an increase in the production and secretion of triglyceride-rich very low density lipoprotein (VLDL-TG) [54–56]. Taken together, this suggests that increased plasma TG concentrations may represent an adaptive response to hepatic insulin resistance. In the present study, we observed a reduction in insulin sensitivity but no change in fasting TG. This finding is in line with previous short-term overfeeding studies, which have observed impairments in glycemic control/reductions in insulin sensitivity alongside unchanged or even reduced fasting TG concentrations [24,25,28]. We did, however, observe a significant trial \times time interaction for plasma TG across the 2-h OGTT, which appears to be due to a more dynamic/temporal postprandial response to glucose ingestion after overfeeding. Previously, it has been shown that there exists a TG storage pool within the enterocytes in which a proportion of meal-derived fatty acids are stored [57]. The amount of TG stored in this pool is related to the amount of dietary fat consumed [58]. These TGs are mobilized and secreted into the circulation following subsequent feeding, a response which has been termed “the second meal effect” [59–61]. Therefore, the increase in plasma TG concentration we observed following glucose ingestion after the single day of high-fat overfeeding is likely to be attributable to the second meal effect due to a larger amount of dietary TG being stored during the intervention period.

The cohort we studied were young, healthy, non-obese males and females who were recreationally active; it is therefore likely that they were relatively metabolically flexible, and somewhat able to adapt to the 1-day diet challenge. Indeed, while we saw significant increases in the postprandial glucose

response after overfeeding, circulating glucose levels at 2 h post-glucose load were considerably lower than the diagnostic values of impaired glucose tolerance (i.e., 7.7–11 mmol/L) [62]. It is plausible that the dietary intervention used in the current study may produce a more dramatic effect in populations at risk of developing T2DM (e.g., sedentary, overweight individuals).

5. Conclusions

In conclusion, our pilot data strongly suggest that replicating excessive binge eating behavior through a single day of high-fat overfeeding is sufficient to impair whole-body insulin sensitivity in young, healthy individuals. Further research is required to elucidate the mechanisms underpinning this response and establish whether these effects persist after returning to normal eating behavior and/or whether repeated periods of binge eating leads to a progressive worsening of glycemic control. Based on our data, it is plausible to suggest that the metabolic effects of binge eating may have more marked effects in individuals at risk of insulin resistance or the metabolic syndrome.

Acknowledgments: The cost of consumables and analysis was covered by the lead investigators own institutional research budget.

Author Contributions: S.A.P. assisted with the design of the study, collected the data and wrote the manuscript. R.M.W. performed dietary analysis, collected the data and assisted with the preparation of the manuscript. L.H. assisted with the preparation of the manuscript. C.J.H. designed the study, collected the data and co-wrote the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Table A1. Example diet plan for one of the subjects.

Breakfast	
Foods	Pork sausages (230 g), streaky bacon (120 g), fried eggs (180 g), fried white bread (36 g), whole milk (300 mL)
Protein (g)	86
Carbohydrate (g)	52
Fat (g)	127
Energy (kJ)	7045
% of the days intake	26
Lunch	
Foods	White bread (72 g), butter (15 g), cheddar cheese (70 g), mayonnaise (20 g), sausage roll (90 g)
Protein (g)	31
Carbohydrate (g)	65
Fat (g)	86
Energy (kJ)	4814
% of the days intake	17
Snack	
Foods	Pork pie (200 g)
Protein (g)	22
Carbohydrate (g)	47
Fat (g)	51
Energy (kJ)	3060
% of the days intake	11

Table A1. Cont.

Dinner	
Foods	Beef burgers (300 g), streaky bacon (120 g), cheddar cheese (90 g), coleslaw (150 g)
Protein (g)	95
Carbohydrate (g)	7
Fat (g)	173
Energy (kJ)	8135
% of the days intake	30
Dessert	
Foods	Chocolate chip muffin (70 g), double cream (150 mL)
Protein (g)	6
Carbohydrate (g)	37
Fat (g)	98
Energy (kJ)	4357
% of the days intake	16
Total intake	
Protein (g)	240
Carbohydrate (g)	209
Fat (g)	535
Energy (kJ)	27,411

Water intake was allowed ad libitum throughout the day.

References

- Brownley, K.A.; Berkman, N.D.; Sedway, J.A.; Lohr, K.N.; Bulik, C.M. Binge eating disorder treatment: A systematic review of randomized controlled trials. *Int. J. Eat. Disord.* **2007**, *40*, 337–348. [[CrossRef](#)] [[PubMed](#)]
- Davis, C.; Levitan, R.D.; Carter, J.; Kaplan, A.S.; Reid, C.; Curtis, C.; Patte, K.; Kennedy, J.L. Personality and eating behaviors: A case-control study of binge eating disorder. *Int. J. Eat. Disord.* **2008**, *41*, 243–250. [[CrossRef](#)] [[PubMed](#)]
- Abraham, T.M.; Massaro, J.M.; Hoffmann, U.; Yanovski, J.A.; Fox, C.S. Metabolic characterization of adults with binge eating in the general population: The framingham heart study. *Obesity* **2014**, *22*, 2441–2449. [[CrossRef](#)] [[PubMed](#)]
- De Zwaan, M. Binge eating disorder and obesity. *Int. J. Obes. Relat. Metab. Disord.* **2001**, *25*, S51–S55. [[CrossRef](#)] [[PubMed](#)]
- Hudson, J.I.; Pope, H.G., Jr. Genetic epidemiology of eating disorders and co-occurring conditions: The role of endophenotypes. *Int. J. Eat. Disord.* **2007**, *40*, S76–S78. [[CrossRef](#)]
- Bray, G.A. Medical consequences of obesity. *J. Clin. Endocrinol. Metab.* **2004**, *89*, 2583–2589. [[CrossRef](#)] [[PubMed](#)]
- Mathes, W.F.; Brownley, K.A.; Mo, X.; Bulik, C.M. The biology of binge eating. *Appetite* **2009**, *52*, 545–553. [[CrossRef](#)] [[PubMed](#)]
- Heatherton, T.F.; Baumeister, R.F. Binge eating as escape from self-awareness. *Psychol. Bull.* **1991**, *110*, 86–108. [[CrossRef](#)] [[PubMed](#)]
- O'Connor, D.B.; Jones, F.; Conner, M.; McMillan, B.; Ferguson, E. Effects of daily hassles and eating style on eating behavior. *Health Psychol.* **2008**, *27*, S20–S31. [[CrossRef](#)] [[PubMed](#)]
- Deer, J.; Koska, J.; Ozias, M.; Reaven, P. Dietary models of insulin resistance. *Metabolism* **2015**, *64*, 163–171. [[CrossRef](#)] [[PubMed](#)]
- Yki-Jarvinen, H. Nutritional modulation of non-alcoholic fatty liver disease and insulin resistance. *Nutrients* **2015**, *7*, 9127–9138. [[CrossRef](#)] [[PubMed](#)]
- Vessby, B.; Uusitupa, M.; Hermansen, K.; Riccardi, G.; Rivellese, A.A.; Tapsell, L.C.; Nalsen, C.; Berglund, L.; Louheranta, A.; Rasmussen, B.M.; et al. Substituting dietary saturated for monounsaturated fat impairs insulin sensitivity in healthy men and women: The kanwu study. *Diabetologia* **2001**, *44*, 312–319. [[CrossRef](#)] [[PubMed](#)]

13. Riccardi, G.; Giacco, R.; Rivellese, A.A. Dietary fat, insulin sensitivity and the metabolic syndrome. *Clin. Nutr.* **2004**, *23*, 447–456. [[CrossRef](#)] [[PubMed](#)]
14. Eckel, R.H.; Grundy, S.M.; Zimmet, P.Z. The metabolic syndrome. *Lancet* **2005**, *365*, 1415–1428. [[CrossRef](#)]
15. Gordon, E.S. Non-esterified fatty acids in the blood of obese and lean subjects. *Am. J. Clin. Nutr.* **1960**, *8*, 740–747.
16. Opie, L.H.; Walfish, P.G. Plasma free fatty acid concentrations in obesity. *N. Engl. J. Med.* **1963**, *268*, 757–760. [[CrossRef](#)] [[PubMed](#)]
17. Karpe, F.; Dickmann, J.R.; Frayn, K.N. Fatty acids, obesity, and insulin resistance: Time for a reevaluation. *Diabetes* **2011**, *60*, 2441–2449. [[CrossRef](#)] [[PubMed](#)]
18. Boden, G.; Jadali, F. Effects of lipid on basal carbohydrate metabolism in normal men. *Diabetes* **1991**, *40*, 686–692. [[CrossRef](#)] [[PubMed](#)]
19. Boden, G.; Jadali, F.; White, J.; Liang, Y.; Mozzoli, M.; Chen, X.; Coleman, E.; Smith, C. Effects of fat on insulin-stimulated carbohydrate metabolism in normal men. *J. Clin. Investig.* **1991**, *88*, 960–966. [[CrossRef](#)] [[PubMed](#)]
20. Boden, G.; She, P.; Mozzoli, M.; Cheung, P.; Gumireddy, K.; Reddy, P.; Xiang, X.; Luo, Z.; Ruderman, N. Free fatty acids produce insulin resistance and activate the proinflammatory nuclear factor-kappaB pathway in rat liver. *Diabetes* **2005**, *54*, 3458–3465. [[CrossRef](#)] [[PubMed](#)]
21. Reeds, D.N.; Stuart, C.A.; Perez, O.; Klein, S. Adipose tissue, hepatic, and skeletal muscle insulin sensitivity in extremely obese subjects with acanthosis nigricans. *Metabolism* **2006**, *55*, 1658–1663. [[CrossRef](#)] [[PubMed](#)]
22. Reaven, G.M.; Hollenbeck, C.; Jeng, C.Y.; Wu, M.S.; Chen, Y.D. Measurement of plasma glucose, free fatty acid, lactate, and insulin for 24 h in patients with niddm. *Diabetes* **1988**, *37*, 1020–1024. [[CrossRef](#)] [[PubMed](#)]
23. Lagerpusch, M.; Bosy-Westphal, A.; Kehden, B.; Peters, A.; Muller, M.J. Effects of brief perturbations in energy balance on indices of glucose homeostasis in healthy lean men. *Int. J. Obes. (Lond)*. **2012**, *36*, 1094–1101. [[CrossRef](#)] [[PubMed](#)]
24. Parry, S.A.; Smith, J.R.; Corbett, T.R.; Woods, R.M.; Hulston, C.J. Short-term, high-fat overfeeding impairs glycaemic control but does not alter gut hormone responses to a mixed meal tolerance test in healthy, normal-weight individuals. *Br. J. Nutr.* **2017**, *117*, 48–55. [[CrossRef](#)] [[PubMed](#)]
25. Wulan, S.N.; Westerterp, K.R.; Plasqui, G. Metabolic profile before and after short-term overfeeding with a high-fat diet: A comparison between south asian and white men. *Br. J. Nutr.* **2014**, *111*, 1853–1861. [[CrossRef](#)] [[PubMed](#)]
26. Brons, C.; Jensen, C.B.; Storgaard, H.; Hiscock, N.J.; White, A.; Appel, J.S.; Jacobsen, S.; Nilsson, E.; Larsen, C.M.; Astrup, A.; et al. Impact of short-term high-fat feeding on glucose and insulin metabolism in young healthy men. *J. Physiol.* **2009**, *587*, 2387–2397. [[CrossRef](#)] [[PubMed](#)]
27. Cornford, A.S.; Hinko, A.; Nelson, R.K.; Barkan, A.L.; Horowitz, J.F. Rapid development of systemic insulin resistance with overeating is not accompanied by robust changes in skeletal muscle glucose and lipid metabolism. *Appl. Physiol. Nutr. Metab.* **2013**, *38*, 512–519. [[CrossRef](#)] [[PubMed](#)]
28. Hulston, C.J.; Churnside, A.A.; Venables, M.C. Probiotic supplementation prevents high-fat, overfeeding-induced insulin resistance in human subjects. *Br. J. Nutr.* **2015**, *113*, 596–602. [[CrossRef](#)] [[PubMed](#)]
29. Nowotny, B.; Zahiragic, L.; Krog, D.; Nowotny, P.J.; Herder, C.; Carstensen, M.; Yoshimura, T.; Szendroedi, J.; Phielix, E.; Schadewaldt, P.; et al. Mechanisms underlying the onset of oral lipid-induced skeletal muscle insulin resistance in humans. *Diabetes* **2013**, *62*, 2240–2248. [[CrossRef](#)] [[PubMed](#)]
30. Mifflin, M.D.; St Jeor, S.T.; Hill, L.A.; Scott, B.J.; Daugherty, S.A.; Koh, Y.O. A new predictive equation for resting energy expenditure in healthy individuals. *Am. J. Clin. Nutr.* **1990**, *51*, 241–247. [[PubMed](#)]
31. Matsuda, M.; DeFronzo, R.A. Insulin sensitivity indices obtained from oral glucose tolerance testing: Comparison with the euglycemic insulin clamp. *Diabetes Care* **1999**, *22*, 1462–1470. [[CrossRef](#)] [[PubMed](#)]
32. Rogers, P.J.; Hill, A.J. Breakdown of dietary restraint following mere exposure to food stimuli: Interrelationships between restraint, hunger, salivation, and food intake. *Addict. Behav.* **1989**, *14*, 387–397. [[CrossRef](#)]
33. Waters, A.; Hill, A.; Waller, G. Internal and external antecedents of binge eating episodes in a group of women with bulimia nervosa. *Int. J. Eat. Disord.* **2001**, *29*, 17–22. [[CrossRef](#)]
34. Adochio, R.L.; Leitner, J.W.; Gray, K.; Draznin, B.; Cornier, M.A. Early responses of insulin signaling to high-carbohydrate and high-fat overfeeding. *Nutr. Metab. (Lond)*. **2009**, *6*, 37. [[CrossRef](#)] [[PubMed](#)]

35. Ferrannini, E.; Bjorkman, O.; Reichard, G.A., Jr.; Pilo, A.; Olsson, M.; Wahren, J.; DeFronzo, R.A. The disposal of an oral glucose load in healthy subjects: A quantitative study. *Diabetes* **1985**, *34*, 580–588. [[CrossRef](#)] [[PubMed](#)]
36. Woerle, H.J.; Meyer, C.; Dostou, J.M.; Gosmanov, N.R.; Islam, N.; Popa, E.; Wittlin, S.D.; Welle, S.L.; Gerich, J.E. Pathways for glucose disposal after meal ingestion in humans. *Am. J. Physiol. Endocrinol. Metab.* **2003**, *284*, E716–E725. [[CrossRef](#)] [[PubMed](#)]
37. Bell, P.M.; Firth, R.G.; Rizza, R.A. Assessment of the postprandial pattern of glucose metabolism in nondiabetic subjects and patients with non-insulin-dependent diabetes mellitus using a simultaneous infusion of [2(3)h] and [3(3)h] glucose. *Metabolism* **1989**, *38*, 38–45. [[CrossRef](#)]
38. Butler, P.C.; Rizza, R.A. Contribution to postprandial hyperglycemia and effect on initial splanchnic glucose clearance of hepatic glucose cycling in glucose-intolerant or niddm patients. *Diabetes* **1991**, *40*, 73–81. [[CrossRef](#)] [[PubMed](#)]
39. Ferrannini, E.; Simonson, D.C.; Katz, L.D.; Reichard, G., Jr.; Bevilacqua, S.; Barrett, E.J.; Olsson, M.; DeFronzo, R.A. The disposal of an oral glucose load in patients with non-insulin-dependent diabetes. *Metabolism* **1988**, *37*, 79–85. [[CrossRef](#)]
40. Ferrannini, E.; Wahren, J.; Felig, P.; DeFronzo, R.A. The role of fractional glucose extraction in the regulation of splanchnic glucose metabolism in normal and diabetic man. *Metabolism* **1980**, *29*, 28–35. [[CrossRef](#)]
41. Firth, R.G.; Bell, P.M.; Marsh, H.M.; Hansen, I.; Rizza, R.A. Postprandial hyperglycemia in patients with noninsulin-dependent diabetes mellitus. Role of hepatic and extrahepatic tissues. *J. Clin. Investig.* **1986**, *77*, 1525–1532. [[CrossRef](#)] [[PubMed](#)]
42. McMahan, M.; Marsh, H.M.; Rizza, R.A. Effects of basal insulin supplementation on disposition of mixed meal in obese patients with niddm. *Diabetes* **1989**, *38*, 291–303. [[CrossRef](#)] [[PubMed](#)]
43. Meyer, C.; Woerle, H.J.; Dostou, J.M.; Welle, S.L.; Gerich, J.E. Abnormal renal, hepatic, and muscle glucose metabolism following glucose ingestion in type 2 diabetes. *Am. J. Physiol. Endocrinol. Metab.* **2004**, *287*, E1049–E1056. [[CrossRef](#)] [[PubMed](#)]
44. Woerle, H.J.; Szoke, E.; Meyer, C.; Dostou, J.M.; Wittlin, S.D.; Gosmanov, N.R.; Welle, S.L.; Gerich, J.E. Mechanisms for abnormal postprandial glucose metabolism in type 2 diabetes. *Am. J. Physiol. Endocrinol. Metab.* **2006**, *290*, E67–E77. [[CrossRef](#)] [[PubMed](#)]
45. Bartoli, E.; Fra, G.P.; Carnevale Schianca, G.P. The oral glucose tolerance test (ogtt) revisited. *Eur. J. Intern. Med.* **2011**, *22*, 8–12. [[CrossRef](#)] [[PubMed](#)]
46. Stumvoll, M.; Van Haeften, T.; Fritsche, A.; Gerich, J. Oral glucose tolerance test indexes for insulin sensitivity and secretion based on various availabilities of sampling times. *Diabetes Care* **2001**, *24*, 796–797. [[CrossRef](#)] [[PubMed](#)]
47. DeFronzo, R.A.; Tobin, J.D.; Andres, R. Glucose clamp technique: A method for quantifying insulin secretion and resistance. *Am. J. Physiol.* **1979**, *237*, E214–E223. [[PubMed](#)]
48. Finegood, D.T.; Bergman, R.N.; Vranic, M. Estimation of endogenous glucose production during hyperinsulinemic-euglycemic glucose clamps: Comparison of unlabeled and labeled exogenous glucose infusates. *Diabetes* **1987**, *36*, 914–924. [[CrossRef](#)] [[PubMed](#)]
49. Hother-Nielsen, O.; Henriksen, J.E.; Holst, J.J.; Beck-Nielsen, H. Effects of insulin on glucose turnover rates in vivo: Isotope dilution versus constant specific activity technique. *Metabolism* **1996**, *45*, 82–91. [[CrossRef](#)]
50. Knudsen, S.H.; Hansen, L.S.; Pedersen, M.; Dejgaard, T.; Hansen, J.; Hall, G.V.; Thomsen, C.; Solomon, T.P.; Pedersen, B.K.; Krogh-Madsen, R. Changes in insulin sensitivity precede changes in body composition during 14 days of step reduction combined with overfeeding in healthy young men. *J. Appl. Physiol. (1985)* **2012**, *113*, 7–15. [[CrossRef](#)] [[PubMed](#)]
51. Zick, Y. Ser/Thr phosphorylation of IRS proteins: A molecular basis for insulin resistance. *Sci STKE* **2005**, *2005*, pe4. [[CrossRef](#)] [[PubMed](#)]
52. Ginsberg, H.N.; Zhang, Y.L.; Hernandez-Ono, A. Regulation of plasma triglycerides in insulin resistance and diabetes. *Arch. Med. Res.* **2005**, *36*, 232–240. [[CrossRef](#)] [[PubMed](#)]
53. Pramfalk, C.; Pavlides, M.; Banerjee, R.; McNeil, C.A.; Neubauer, S.; Karpe, F.; Hodson, L. Fasting plasma insulin concentrations are associated with changes in hepatic fatty acid synthesis and partitioning prior to changes in liver fat content in healthy adults. *Diabetes* **2016**, *65*, 1858–1867. [[CrossRef](#)] [[PubMed](#)]

54. Hodson, L.; Banerjee, R.; Rial, B.; Arlt, W.; Adiels, M.; Boren, J.; Marinou, K.; Fisher, C.; Mostad, I.L.; Stratton, I.M.; et al. Menopausal status and abdominal obesity are significant determinants of hepatic lipid metabolism in women. *J. Am. Heart Assoc.* **2015**, *4*, e002258. [[CrossRef](#)] [[PubMed](#)]
55. Lambert, J.E.; Parks, E.J. Postprandial metabolism of meal triglyceride in humans. *Biochim. Biophys. Acta* **2012**, *1821*, 721–726. [[CrossRef](#)] [[PubMed](#)]
56. Matikainen, N.; Adiels, M.; Soderlund, S.; Stenabb, S.; Ahola, T.; Hakkarainen, A.; Boren, J.; Taskinen, M.R. Hepatic lipogenesis and a marker of hepatic lipid oxidation, predict postprandial responses of triglyceride-rich lipoproteins. *Obesity* **2014**, *22*, 1854–1859. [[CrossRef](#)] [[PubMed](#)]
57. Robertson, M.D.; Parkes, M.; Warren, B.F.; Ferguson, D.J.; Jackson, K.G.; Jewell, D.P.; Frayn, K.N. Mobilisation of enterocyte fat stores by oral glucose in humans. *Gut* **2003**, *52*, 834–839. [[CrossRef](#)] [[PubMed](#)]
58. Zhu, J.; Lee, B.; Buhman, K.K.; Cheng, J.X. A dynamic, cytoplasmic triacylglycerol pool in enterocytes revealed by ex vivo and in vivo coherent anti-stokes raman scattering imaging. *J. Lipid Res.* **2009**, *50*, 1080–1089. [[CrossRef](#)] [[PubMed](#)]
59. Fielding, B.A.; Callow, J.; Owen, R.M.; Samra, J.S.; Matthews, D.R.; Frayn, K.N. Postprandial lipemia: The origin of an early peak studied by specific dietary fatty acid intake during sequential meals. *Am. J. Clin. Nutr.* **1996**, *63*, 36–41. [[PubMed](#)]
60. Jackson, K.G.; Robertson, M.D.; Fielding, B.A.; Frayn, K.N.; Williams, C.M. Olive oil increases the number of triacylglycerol-rich chylomicron particles compared with other oils: An effect retained when a second standard meal is fed. *Am. J. Clin. Nutr.* **2002**, *76*, 942–949. [[PubMed](#)]
61. Robertson, M.D.; Henderson, R.A.; Vist, G.E.; Rumsey, R.D. Extended effects of evening meal carbohydrate-to-fat ratio on fasting and postprandial substrate metabolism. *Am. J. Clin. Nutr.* **2002**, *75*, 505–510. [[PubMed](#)]
62. Nathan, D.M.; Davidson, M.B.; DeFronzo, R.A.; Heine, R.J.; Henry, R.R.; Pratley, R.; Zinman, B.; American Diabetes, A. Impaired fasting glucose and impaired glucose tolerance: Implications for care. *Diabetes Care* **2007**, *30*, 753–759. [[CrossRef](#)] [[PubMed](#)]



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).