

## Loughborough University Institutional Repository

---

### *Erratum: colossal magneto-optical conductivity in doped manganites*

This item was submitted to Loughborough University's Institutional Repository by the/an author.

**Citation:** ALEXANDROV, A.S. and BRATKOVSKY, A.M., 2000. Erratum: colossal magneto-optical conductivity in doped manganites. *Physical Review B*, 61 (5), pp. 3748

**Additional Information:**

- This article was published in the journal *Physical Review B* [© American Physical Society]. The definitive version is available at: <http://link.aps.org/doi/10.1103/PhysRevB.61.3748>

**Metadata Record:** <https://dspace.lboro.ac.uk/2134/4281>

**Version:** Published

**Publisher:** © American Physical Society

Please cite the published version.

This item was submitted to Loughborough's Institutional Repository (<https://dspace.lboro.ac.uk/>) by the author and is made available under the following Creative Commons Licence conditions.



For the full text of this licence, please go to:  
<http://creativecommons.org/licenses/by-nc-nd/2.5/>

---

---

**ERRATA**

---

---

**Erratum: Colossal magneto-optical conductivity in doped manganites**  
**[Phys. Rev. B 60, 6215 (1999)]**

A. S. Alexandrov and A. M. Bratkovsky

There is a typo in the comment following Eq. (9) for  $n_c(h)$  close to the ferromagnetic phase transition on the paramagnetic side. It should read:

Here we assume that  $w \gg t_c$  (in the opposite limit,  $w \ll t_c$ , the preexponential factor will be 2 instead of  $t_c/w$ ). The factors result from assuming that the total number of states in the bipolaron (impurity) band equals the doping level  $x$ . If we were to normalize to unity instead of  $x$ , as in Ref. 1, we would obtain the prefactors  $(t_c/w)\sqrt{x/2}$  and  $\sqrt{2x}$ , respectively.

This correction has no effect on our results and discussion.

<sup>1</sup>A. S. Alexandrov and A. M. Bratkovsky, Phys. Rev. Lett. **82**, 141 (1999).